



















Protected areas: A focus on Brazilian freshwater biodiversity

Valter M. Azevedo-Santos¹  | Renata G. Frederico²  | Camila K. Fagundes³ |
 Paulo S. Pompeu⁴  | Fernando M. Pelicice⁵  | André A. Padial^{6,7}  |
 Marcos G. Nogueira¹ | Philip M. Fearnside⁸  | Luciano B. Lima^{9,10}  |
 Vanessa S. Daga¹¹  | Fagner J. M. Oliveira^{9,12}  | Jean R. S. Vitule^{7,11}  |
 Marcos Callisto¹³  | Angelo A. Agostinho¹²  | Francisco A. Esteves¹⁴ |
 Dilermando P. Lima-Junior⁹ | André L. B. Magalhães¹⁵  | José Sabino¹⁶  |
 Roger P. Mormul¹²  | Daniel Grasel¹⁷  | Jansen Zuanon⁸  | Fábio S. Vilella¹⁸ |
 Raoul Henry¹ 

¹Universidade Estadual Paulista "Júlio de Mesquita Filho", Botucatu, São Paulo, Brazil

²Instituto de Ciências Biológicas, Universidade Federal do Pará, Belém, Pará, Brazil

³Wildlife Conservation Society, Brazil Program, Manaus, Amazonas, Brazil

⁴Laboratório de Ecologia de Peixes, Departamento de Biologia, Universidade Federal de Lavras, Lavras, Minas Gerais, Brazil

⁵Núcleo de Estudos Ambientais, Universidade Federal de Tocantins, Porto Nacional, Tocantins, Brazil

⁶Laboratório de Análise e Síntese em Biodiversidade (LASB), Departamento de Botânica, Universidade Federal do Paraná, Curitiba, Paraná, Brazil

⁷Programa de Pós-Graduação em Ecologia e Conservação, Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba, Paraná, Brazil

⁸Instituto Nacional de Pesquisas da Amazônia (INPA), Manaus, Amazonas, Brazil

⁹Laboratório de Ecologia e Conservação de Ecossistemas Aquáticos, Universidade Federal de Mato Grosso, Pontal do Araguaia, Mato Grosso, Brazil

¹⁰Programa de Pós-Graduação em Ecologia e Conservação, Universidade do Estado de Mato Grosso, Nova Xavantina, Mato Grosso, Brazil

¹¹Laboratório de Ecologia e Conservação (LEC), Departamento de Engenharia Ambiental, Setor de Tecnologia, Universidade Federal do Paraná, Curitiba, Paraná, Brazil

¹²Programa de Pós-Graduação em Ecologia de Ambientes Aquáticos Continentais, Universidade Estadual de Maringá, Maringá, Paraná, Brazil

¹³Laboratório de Ecologia de Bentos, Departamento de Genética, Ecologia e Evolução, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Belo Horizonte, Minas Gerais, Brazil

¹⁴Núcleo em Ecologia e Desenvolvimento Sócio Ambiental de Macaé, Universidade Federal do Rio de Janeiro, Macaé, Rio de Janeiro, Brazil

¹⁵Programa de Pós-Graduação em Tecnologias para o Desenvolvimento Sustentável, Universidade Federal de São João Del Rei, Ouro Branco, Minas Gerais, Brazil

¹⁶Universidade Anhangüera - Uniderp, Campo Grande, Mato Grosso do Sul, Brazil

¹⁷Programa de Pós-Graduação em Botânica, Instituto de Biotecnologia, Universidade Federal do Rio Grande do Sul, Porto Alegre, Rio Grande do Sul, Brazil

¹⁸Simbiota Consultoria Ambiental Ltda., Venâncio Aires, Rio Grande do Sul, Brazil

Correspondence

Valter M. Azevedo-Santos, Universidade Estadual Paulista "Júlio de Mesquita Filho", Botucatu, São Paulo, Brazil.
 Email: valter.ecologia@gmail.com

Funding information

Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES); CNPq/FAPESPA, Grant/Award Number:

Abstract

Brazil has a variety of aquatic ecosystems and rich freshwater biodiversity, but these components have been constantly damaged by the expansion of unsustainable activities. An array of different conservation strategies is needed, especially the creation of protected areas (PAs, hereafter). However, Brazil's PAs are biased towards terrestrial ecosystems and we argue that current PAs have limited efficacy in the protection of

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2018 The Authors. *Diversity and Distributions* Published by John Wiley & Sons Ltd

ICAAF 094/2016; Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq); Fundação de Amparo à Pesquisa do Estado de Mato Grosso (FAPEMAT); Programa de Pesquisa e Desenvolvimento Tecnológico do Setor de Energia Elétrica- Companhia Energética de Minas Gerais

Editor: Robert Cowie

freshwater biodiversity. New PAs should better consider aquatic environments, covering entire basins, rivers and other freshwater habitats. We recommend ways to implement these PAs and provide guidance to avoid social impacts. Freshwater systems in Brazil provide essential goods and services but these ecosystems are being rapidly degraded and will be lost if not adequately protected.

KEYWORDS

conservation units, human impacts, indigenous land, inland waters, Neotropical region

1 | INTRODUCTION

Brazil has a large number of freshwater ecosystems, many of which have high biodiversity (Abell et al., 2008; Agostinho, Thomaz, & Gomes, 2005; Turtle Taxonomy Working Group, 2017) and endemism (e.g., Bertaco & Carvalho, 2010; Daga, Debona, Abilhoa, Gubiani, & Vitule, 2016). However, many of these ecosystems have been disrupted by human activities, threatening the maintenance of native biodiversity and ecosystem services. This scenario points to the urgent need for a variety of different conservation strategies in the scientific, educational and political realms (Azevedo-Santos et al., 2017; Karam-Gemael, Loyola, Penha, & Izzo, 2018; Pelicice et al., 2017).

Creation of protected areas (PAs, hereafter) in different regions of Brazil represented an advance for conservation policies (Mittermeier, Fonseca, Rylands, & Brandon, 2005), especially because they can prevent deforestation (Nolte, Agrawal, Silvius, & Soares-Filho, 2013). However, current PAs have limited efficacy in protecting freshwater biodiversity (e.g., Fagundes, Vogt, & De Marco, 2016; Frederico, Zuanon, & De Marco, 2018). The creation of PAs in Brazil was strongly based on terrestrial biota and the threats to them, in general ignoring aquatic species and ecosystems (Fagundes et al., 2016; Frederico et al., 2018).

In this short essay, we question the efficacy of Brazilian PAs (only Conservation Units) in protecting freshwater environments and associated biodiversity. We argue that new PAs, based on the current categories of the National System of Conservation Units (SNUC, 2000), should explicitly incorporate freshwater biodiversity. In addition, we provide some guidance for avoiding social conflicts that may emerge from the establishment of these new PAs. Finally, we stress the urgency of creation of new PAs in the face of the ongoing expansion of unsustainable activities and the progressive loss of Brazilian freshwater ecosystems and biodiversity.

2 | FRESHWATER BIODIVERSITY UNPROTECTED

Most Brazilian PAs were created to protect terrestrial biodiversity (mainly vegetation cover, and strongly focused on forests; Overbeck et al., 2015). In the Brazilian Amazon, approximately 60 million hectares are protected but studies indicate that PAs have limited value

for freshwater chelonians (Fagundes et al., 2016) and stream fishes (Frederico et al., 2018). In addition, the decline of large migratory fishes (e.g., Petrere, Barthem, Cordoba, & Gomez, 2004; Castello et al., 2013; Correa et al., 2015) indicates that existing PAs are not enough to maintain population dynamics and fisheries stocks. These fishes demand large areas to complete their life cycles, posing a challenge to conservation policies. Amazon PAs are, in general, vast ("mega-reserves", sensu Peres, 2005), but covering large forested areas does not automatically protect their freshwater biodiversity (Fagundes et al., 2016; Frederico et al., 2018).

In Brazil's southeast region, PAs are also biased in favor of protecting terrestrial biodiversity. In this region, where different biomes are found but most of its native vegetation has been removed, PAs "typically comprise hilltop areas and therefore exclude a large number of species that are not present in this type of environment" (Pompeu, Reis, Gandini, Souza, & Favero, 2009:665). Such a scenario excludes most freshwater habitats and their biota, with the possible exception of the smallest headwater streams. This situation does not necessarily apply to all regions of the country (e.g., Amazonia), but it represents another failure of the current conservation strategies in Brazil.

From a freshwater perspective, the most important limitation is that existing PAs, in general, cover only short stretches of river systems (Figure 1) or, in some cases, a small fraction of their headwaters. This pattern conflicts with the nature of fluvial networks, which are dynamic hierarchical macro-systems that strongly depend on connectivity to maintain their ecological integrity (Grant, Lowe, & Fagan, 2007; McCluney et al., 2014). Distant locations (i.e., habitat patches) are connected by fluvial watercourses, in which species disperse and occupy different habitats throughout their life cycles. In addition, local disturbances may propagate along the watercourses, especially in the downstream direction. This means that human actions in a river basin can spread through the system and affect any PAs that have been established (Saunders, Meeuwig, & Vincent, 2002), making the entire system vulnerable to anthropogenic actions occurring at different locations (e.g., pollution, mining, overfishing, damming). PAs that cover only segments of a river cannot secure the integrity, heterogeneity and dynamism of fluvial networks, which potentially threaten most freshwater organisms.

Restricting the protection to river segments does not allow maintenance of system functionality and complex biological dynamics. Migratory species, including some fishes, have complex




FIGURE 1 Protected areas (Conservation Units) and main Brazilian rivers. The shapefile “Conservation Units (all)” (MMA, 2018) and the shapefile “Principal Rivers” (sensu ANA, 2018) were used to the map, which was constructed using Quantum GIS (QGIS Development Team, 2018)

life cycles that demand spatial heterogeneity, that is, a diversity of habitats and specific conditions (e.g., Carolsfeld, Harvey, Ross, & Baer, 2003). The natural flow regime is another important function that single PAs cannot preserve because flow dynamics depend on a number of variables acting at the regional scale (Humphries, Keckeis, & Finlayson, 2014). For the Amazon basin, Castello et al. (2013:223) argued that “The protected area network provides protection against overharvesting and riparian deforestation, but does not protect freshwater ecosystems from the far-reaching impacts of dams”. Impoundments have such effects in both upstream and downstream directions (see Agostinho, Pelicice, & Gomes, 2008), affecting the biodiversity in PAs. The invasion of non-native species is another challenge, because human activity

(e.g., aquaculture) outside PAs introduces species that may reach PAs (Figure 2).

3 | PROTECTED AREAS FOR FRESHWATER BIODIVERSITY

3.1 | Focusing on freshwaters

The National System of Conservation Units (“SNUC” in Portuguese) set the guidelines for most of the country’s PAs (known as “Conservation Units” [SNUC, 2000]). In principle, PAs in all categories of the SNUC could be established to protect freshwater biodiversity. For instance, the category “Wildlife Refuge” (*Refúgio de Vida*




FIGURE 2 Hypothetical illustration showing how local PAs are inefficient for protecting freshwater biodiversity in river systems that are subjected to several human disturbances

Silvestre) could be used to preserve critical habitats for migratory species (e.g., spawning and nursery grounds) and other areas with essential functions for aquatic species (e.g., ponds, lakes and floodplains). A detailed analysis is needed to reveal how each PA category applies to different freshwater ecosystems, as well as the need for new categories. Here we just explore two approaches that may initiate the discussion about freshwater PAs in Brazil, which may be applicable to other countries with high levels of freshwater biodiversity. The first approach should be combined with the second, since they complement each other.

The first approach concerns the features of waterbodies or basins to be protected. We suggest, as a first step, protecting: (a) areas relevant for migratory species (whether threatened or not), (b) areas with endemic or threatened species and/or (c) areas with high biological diversity (taxonomic, functional and/or phylogenetic). Systematic conservation planning can indicate biodiversity gaps and priority sites for the development of PAs and other conservation strategies (Fearnside, 2015; Fearnside & Ferraz, 1995; Fonseca & Venticinque, 2018). The Brazilian government has adopted systematic planning (MMA, 2016), but as usual, they are biased towards terrestrial biodiversity. This is a global trend, since there are few studies that propose systematic planning for inland aquatic ecosystems (e.g., Carrizo et al., 2017).

The second approach is to consider the possibility of protecting entire basins (small to large). This measure is more desirable and would take into account all elements of a basin (e.g., watercourses, ponds, lakes, etc), in addition to functional aspects (e.g., the natural

flow regime). If it is not possible to protect an entire basin, at least waterbodies such as rivers should be protected, including their entire length (i.e., headwaters to mouth) and adjacent areas (e.g., wetlands, upland buffers). Brazil has legal mechanisms, not included in the SNUC, to protect rivers; but this strategy has been rarely employed. For instance, law 15.082/2004 creates “permanent preservation rivers” (“*rios de preservação permanente*”) that are valid in the state of Minas Gerais. This law represents a breakthrough in the protection of freshwater environments; it is clearly stated that entire rivers must be protected.

3.2 | Avoiding social impacts of new PAs

To avoid social impacts and conflicts, conservation policies should balance creation of restrictive (i.e., integral protection) and less restrictive (i.e., sustainable use) PAs. Protection of rivers and basins is more challenging than preservation of terrestrial areas. In Brazil and in many other countries, riverine resources sustain basic human needs, including those of indigenous and traditional populations. Implementation of fully PAs would lead to land expropriation and strict use restrictions, causing unacceptable levels of social impact. The severe social impacts of removing traditional people and of destroying their livelihoods are illustrated by recent hydroelectric dams on the Madeira (Fearnside, 2014) and Xingu Rivers (Fearnside, 2017; Magalhães & da Cunha, 2017). The SNUC includes “Integral-Protection Units” (“*Unidades de Proteção Integral*”) and “Sustainable-Use Units” (“*Unidades de Uso Sustentável*”) (SNUC, 2000), but the

second category seems more appropriate for most river basins in Brazil. Artisanal fishing in tropical river systems, for example, is an important activity that tends to be irreplaceable but compatible with less-restrictive PAs. Appropriate management of sustainable-use units is therefore required to combine human activities with the protection of aquatic environments and their ecological functions. Combinations of different PA categories (i.e., geographical mosaics of Conservation Units) must also be evaluated. For example, headwater streams could be protected by integral-protection units, while important downstream reaches could be protected by sustainable-use units.

3.3 | Transboundary freshwater ecosystems

Major South American rivers run across national borders. This is the case of the Paraná, Paraguay and Uruguay rivers in the La Plata basin, which are shared by five countries (Argentina, Brazil, Bolivia, Paraguay and Uruguay). The same is true for several Amazonian tributaries, such as the Madeira, Purus, Solimões (Upper Amazon) and Japurá rivers. These cases will demand international agreements to implement effective PAs, otherwise poor conservation policies in one country will jeopardize protection efforts in another. For example, the ongoing hydropower development in Peru, Colombia and Bolivia (Finer & Jenkins, 2012) will have negative effects on floodplain ecosystems and fish populations in Brazil (Forsberg et al., 2017; Latrubesse et al., 2017; Lees, Peres, Fearnside, Schneider, & Zuanon, 2016). The presence of these macro-river systems in Brazil indicates that conservation efforts will depend on transboundary policies.

4 | THE NEED FOR MORE PAs

Brazil needs more PAs (Overbeck et al., 2015). However, these new areas should also consider freshwater environments and biodiversity because they are subject to a variety of human disturbances (e.g., Alho & Sabino, 2011; Vitule et al., 2015; Garcia, Ribeiro, Roque, Ochoa-Quintero, & Laurance, 2017). For example, a large number of small- and medium-sized dams have been planned or are under construction in Brazil and in other South American countries (Zarfl, Lumsdon, Berlekamp, Tydecks, & Tockner, 2015); hydropower expansion can irreversibly damage biodiversity and ecosystem functioning (Couto & Olden, 2018; Linares, Callisto, & Marques, 2018; Nogueira, Oliveira, & Britto, 2008; Winemiller et al., 2016).

In addition to dams, there are numerous other negative impacts caused by human activities that threaten freshwater biodiversity in Brazil. For instance, non-native freshwater species have been introduced for aquaculture or other activities for decades in the country (e.g., Magalhães et al., 2005; Coelho & Henry, 2017; Pelicice et al., 2017; Coelho, Fernandez, Cesar, Ruocco, & Henry, 2018; Gubiani et al., 2018), which can result in loss of native species and biotic homogenization (Mack et al., 2000; Simberloff et al., 2013; Vitule, Freire, Vazquez, Nuñez, & Simberloff, 2012). Other harmful actions,

such as mining, have high impact on rivers (e.g., Fernandes et al., 2016; Tófoli et al., 2017), and existing PAs will be inefficient in protecting biodiversity from these perturbations.

Lastly we highlight that Brazil is one of the Latin American countries that has agreed to the "Aichi Biodiversity Targets", which list a series of goals for biological conservation worldwide (CBD, 2018). Establishing new PAs for freshwater systems is in accord with this treaty, specifically Target 11 (see CBD, 2018 for more details), and, therefore, Brazil needs to take appropriate measures to create these PAs.

ACKNOWLEDGEMENTS

We are grateful to Robert H. Cowie, for corrections and suggestions, and André Morato, for the Figure 2. V.M.A.S., F.J.M.O., A.L.B.M., and D.G. were supported by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES). R.G.F. was supported by CNPq/FAPESPA (ICAAF 094/2016). P.S.P., F.M.P., P.M.F., M.C. (PQ Process Number: 303380/2015-2), A.A.P. (Process Numbers: 307984/2015-0 and 402828/2016-0), V.S.D. (Process Number: 167382/2017-9), J.R.S.V. (PQ Process Numbers: 310850/2012-6 and 303776/2015-3), J.Z. (PQ Process Number: 313183/2014-7), and A.A.A. were supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). L.B.L. was supported by Fundação de Amparo à Pesquisa do Estado de Mato Grosso (FAPEMAT). M.C. received financial support from Programa de Pesquisa e Desenvolvimento Tecnológico do Setor de Energia Elétrica-Companhia Energética de Minas Gerais (P&D Aneel-Cemig GT-599).

ORCID

Valter M. Azevedo-Santos  <http://orcid.org/0000-0001-8986-6406>

Renata G. Frederico  <https://orcid.org/0000-0002-8858-6426>

Paulo S. Pompeu  <https://orcid.org/0000-0002-7938-1517>

Fernando M. Pelicice  <https://orcid.org/0000-0002-9700-1194>

André A. Padial  <https://orcid.org/0000-0002-8766-5974>

Philip M. Fearnside  <https://orcid.org/0000-0003-3672-9082>

Luciano B. Lima  <http://orcid.org/0000-0002-8286-8035>

Vanessa S. Daga  <http://orcid.org/0000-0003-3707-716X>

Fagner J. M. Oliveira  <https://orcid.org/0000-0003-0602-310X>

Jean R. S. Vitule  <https://orcid.org/0000-0001-6543-7439>

Marcos Callisto  <https://orcid.org/0000-0003-2341-4700>

Angelo A. Agostinho  <https://orcid.org/0000-0002-4707-9444>

André L. B. Magalhães  <https://orcid.org/0000-0002-9463-1836>

José Sabino  <https://orcid.org/0000-0002-6404-5107>

Roger P. Mormul  <https://orcid.org/0000-0001-9020-4784>

Daniel Grasel  <https://orcid.org/0000-0001-6978-7508>

Jansen Zuanon  <https://orcid.org/0000-0001-8354-2750>

Raoul Henry  <http://orcid.org/0000-0002-4000-2524>

REFERENCES

- Abell, R., Thieme, M. L., Revenga, C., Bryer, M., Kottelat, M., Bogutskaya, N., ... Petry, P. (2008). Freshwater ecoregions of the world: A new map of biogeographic units for freshwater biodiversity conservation. *BioScience*, 58, 403–414. <https://doi.org/10.1641/B580507>
- Agência Nacional de Águas (ANA) (2018). *Rios Principais (Principal Rivers)*. Retrieved from <https://metadados.ana.gov.br/geonetwork/srv/pt/main.home?uuxml:id=fb206e35-a86e-44b3-ac3b-5c4d6833c48e>
- Agostinho, A. A., Pelicice, F. M., & Gomes, L. C. (2008). Dams and the fish fauna of the Neotropical region: Impacts and management related to diversity and fisheries. *Brazilian Journal of Biology*, 68, 1119–1132. <https://doi.org/10.1590/S1519-69842008000500019>
- Agostinho, A. A., Thomaz, S. M., & Gomes, L. C. (2005). Conservation of the biodiversity of Brazil's inland waters. *Conservation Biology*, 19, 646–652. <https://doi.org/10.1111/j.1523-1739.2005.00701.x>
- Alho, C. J., & Sabino, J. (2011). A conservation agenda for the Pantanal's biodiversity. *Brazilian Journal of Biology*, 71(Supplement 1), 327–335. <https://doi.org/10.1590/S1519-69842011000200012>
- Azevedo-Santos, V. M., Fearnside, P. M., Oliveira, C. S., Padiá, A. A., Pelicice, F. M., Lima-Junior, D. P., ... Vitule, J. R. S. (2017). Removing the abyss between conservation science and policy decision in Brazil. *Biodiversity and Conservation*, 26, 1745–1752. <https://doi.org/10.1007/s10531-017-1316-x>
- Bertaco, V. A., & Carvalho, F. R. (2010). New species of *Hasemania* (Ostariophysi: Characiformes: Characidae) from Central Brazil, with comments on the endemism of upper rio Tocantins basin, Goiás State. *Neotropical Ichthyology*, 8(1), 27–32.
- Carolsfeld, J., Harvey, B., Ross, C., & Baer, A. (2003). *Migratory fishes of South America: Biology, fisheries and conservation status*. Ottawa, ON: World Fisheries Trust, The World Bank.
- Carrizo, S. F., Lengyel, S., Kapusi, F., Szabolcs, M., Kasperidus, H. D., Scholz, M., ... Darwall, W. (2017). Critical catchments for freshwater biodiversity conservation in Europe: Identification, prioritisation and gap analysis. *Journal of Applied Ecology*, 54, 1209–1218. <https://doi.org/10.1111/1365-2664.12842>
- Castello, L., McGrath, D. G., Hess, L. L., Coe, M. T., Lefebvre, P. A., Petry, P., ... Arantes, C. C. (2013). The vulnerability of Amazon freshwater ecosystems. *Conservation Letters*, 6, 217–222. <https://doi.org/10.1111/conl.12008>
- Coelho, P. N., Fernandez, M. A., Cesar, D. A. S., Ruocco, A. M. C., & Henry, R. (2018). Updated distribution and range expansion of the gastropod invader *Melanoides tuberculata* (Müller, 1774) in Brazilian waters. *BioInvasions Records* (in press).
- Coelho, P. N., & Henry, R. (2017). The small foreigner: New laws will promote the introduction of non-native zooplankton in Brazilian aquatic environments. *Acta Limnologica Brasiliensia*, 29, e7. <https://doi.org/10.1590/s2179-975x0717>
- Convention on Biological Diversity (CBD) (2018). *Aichi Biodiversity Targets*. Retrieved from <https://www.cbd.int/sp/targets/>
- Correa, S. B., Araujo, J. K., Penha, J. M. F., Cunha, C. N., Stevenson, P. R., & Anderson, J. T. (2015). Overfishing disrupts an ancient mutualism between frugivorous fishes and plants in Neotropical wetlands. *Biological Conservation*, 191, 159–167. <https://doi.org/10.1016/j.biocon.2015.06.019>
- Couto, T. B. A., & Olden, J. D. O. (2018). Global proliferation of small hydropower plants – science and policy. *Frontiers in Ecology and the Environment*, 16(2), 91–100. <https://doi.org/10.1002/fee.1746>
- Daga, V. S., Debona, T., Abilhoa, V., Gubiani, É. A., & Vitule, J. R. S. (2016). Non-native fish invasions of a Neotropical ecoregion with high endemism: A review of the Iguazu River. *Aquatic Invasions*, 11, 209–223. <https://doi.org/10.3391/ai.2016.11.2.10>
- Fagundes, C. K., Vogt, R. C., & De Marco, P. Jr (2016). Testing the efficiency of protected areas in the Amazon for conserving freshwater turtles. *Diversity and Distributions*, 22, 123–135. <https://doi.org/10.1111/ddi.12396>
- Fearnside, P. M. (2014). Impacts of Brazil's Madeira River dams: Unlearned lessons for hydroelectric development in Amazonia. *Environmental Science & Policy*, 38, 164–172. <https://doi.org/10.1016/j.envsci.2013.11.004>
- Fearnside, P. M. (2015). Pesquisa sobre conservação na Amazônia brasileira e a sua contribuição para a manutenção da biodiversidade e uso sustentável das florestas tropicais. In I. C. G. Vieira, M. A. G. Jardim, & E. J. P. da Rocha (Eds.), *Amazônia em Tempo: Estudos Climáticos e Socioambientais* (pp. 21–49). Belém, Brazil: Universidade Federal do Pará, Museu Paraense Emílio Goeldi & Embrapa Amazônia Oriental.
- Fearnside, P. M. (2017). Brazil's Belo Monte Dam: Lessons of an Amazonian resource struggle. *Die Erde*, 148(2–3), 167–184. <https://doi.org/10.12854/erde-148-46>
- Fearnside, P. M., & Ferraz, J. (1995). A conservation gap analysis of Brazil's Amazonian vegetation. *Conservation Biology*, 9(5), 1134–1147. <https://doi.org/10.1046/j.1523-1739.1995.9051127.x>
- Fernandes, G. W., Goulart, F. F., Ranieri, B. D., Coelho, M. S., Dales, K., Boesche, N., ... Soares-Filho, B. (2016). Deep into the mud: Ecological and socio-economic impacts of the dam breach in Mariana, Brazil. *Natureza & Conservação*, 14, 35–45. <https://doi.org/10.1016/j.ncon.2016.10.003>
- Finer, M., & Jenkins, C. N. (2012). Proliferation of hydroelectric dams in the Andean Amazon and implications for Andes-Amazon connectivity. *PLoS ONE*, 7(4), e35126. <https://doi.org/10.1371/journal.pone.0035126>
- Fonseca, C. R., & Venticinque, E. M. (2018). Biodiversity conservation gaps in Brazil: A role for systematic conservation planning. *Perspectives in Ecology and Conservation*, 16, 61–67. <https://doi.org/10.1016/j.pecon.2018.03.001>
- Forsberg, B. R., Melack, J. M., Dunne, T., Barthem, R. B., Goulding, M., Paiva, R. C. D., ... Silva, U. L. Jr (2017). The potential impact of new Andean dams on Amazon fluvial ecosystems. *PLoS ONE*, 12(8), e0182254. <https://doi.org/10.1371/journal.pone.0182254>
- Frederico, R. G., Zuanon, J., De Marco, P. Jr (2018). Amazon protected areas and its ability to protect stream-dwelling fish fauna. *Biological Conservation*, 219, 12–19. <https://doi.org/10.1016/j.biocon.2017.12.032>
- Garcia, L. C., Ribeiro, D. B., Roque, F. O., Ochoa-Quintero, J. M., & Laurance, W. F. (2017). Brazil's worst mining disaster: Corporations must be compelled to pay the actual environmental costs. *Ecological Applications*, 27, 5–9. <https://doi.org/10.1002/eap.1461>
- Grant, E. H. C., Lowe, W. H., & Fagan, W. F. (2007). Living in the branches: Population dynamics and ecological processes in dendritic networks. *Ecology Letters*, 10, 165–175. <https://doi.org/10.1111/j.1461-0248.2006.01007.x>
- Gubiani, E. A., Ruaro, R., Ribeiro, V. R., Eichelberger, A. C. A., Bogoni, R. F., Lira, A. D., ... da Graça, W. J. (2018). Non-native fish species in Neotropical freshwaters: How did they arrive, and where did they come from? *Hydrobiologia*, 817, 57–69. <https://doi.org/10.1007/s10750-018-3617-9>
- Humphries, P., Keckeis, H., & Finlayson, B. (2014). The river wave concept: Integrating river ecosystem models. *BioScience*, 64, 870–882. <https://doi.org/10.1093/biosci/biu130>
- Karam-Gemael, M., Loyola, R., Penha, J., & Izzo, T. (2018). Poor alignment of priorities between scientists and policymakers highlights the need for evidence-informed conservation in Brazil. *Perspectives in Ecology and Conservation*, 16, 125–132. <https://doi.org/10.1016/j.pecon.2018.06.002>
- Latrubesse, E. M., Arima, E. Y., Dunne, T., Park, E., Baker, V. R., D'Horta, F. M., ... Stevaux, J. C. (2017). Damming the rivers of the Amazon basin. *Nature*, 546, 363–369. <https://doi.org/10.1038/nature22333>
- Lees, A. C., Peres, C. A., Fearnside, P. M., Schneider, M., & Zuanon, J. A. S. (2016). Hydropower and the future of Amazonian biodiversity.

- Biodiversity and Conservation*, 25, 451–466. <https://doi.org/10.1007/s10531-016-1072-3>
- Linares, M. S., Callisto, M., & Marques, J. C. (2018). Thermodynamic based indicators illustrate how a run-of-river impoundment in neotropical savanna attracts invasive species and alters the benthic macroinvertebrate assemblages' complexity. *Ecological Indicators*, 88, 181–189. <https://doi.org/10.1016/j.ecolind.2018.01.040>
- Mack, R. N., Simberloff, D., Lonsdale, W. M., Evans, H., Clout, M., & Bazzaz, F. A. (2000). Biotic invasions: Causes, epidemiology, global consequences, and control. *Ecological Applications*, 10, 689–710. [https://doi.org/10.1890/1051-0761\(2000\)010\[0689:BICEGC\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2000)010[0689:BICEGC]2.0.CO;2)
- Magalhães, C., Bueno, S. L. S., Bond-Buckup, G., Valenti, W. C., Silva, H. L. M., Kiyohara, F., ... Rocha, S. S. (2005). Exotic species of freshwater decapod crustaceans in the state of São Paulo, Brazil: Records and possible causes of their introduction. *Biodiversity and Conservation*, 14, 1929–1945. <https://doi.org/10.1007/s10531-004-2123-8>
- Magalhães, S. B., & M. C. da Cunha (Eds.) (2017). *A Expulsão de Ribeirinhos em Belo Monte: Relatório da SBPC*. São Paulo, Brazil: Sociedade Brasileira para o Progresso da Ciência-SBPC. Retrieved from <https://portal.sbpnet.org.br/livro/belomonte.pdf>
- McCluney, K. E., Poff, N. L., Palmer, M. A., Thorp, J. H., Poole, G. C., Williams, B. S., ... Baron, J. S. (2014). Riverine macrosystems ecology: Sensitivity, resistance, and resilience of whole river basins with human alterations. *Frontiers in Ecology and the Environment*, 12(1), 48–58. <https://doi.org/10.1890/120367>
- Mittermeier, R. A., Fonseca, G. A. B., Rylands, A. B., & Brandon, K. (2005). Uma breve história da conservação da biodiversidade no Brasil. *Megadiversidade*, 1(1), 14–21.
- MMA (Ministério do Meio Ambiente) (2016). *Caatinga*. Segunda atualização. Retrieved from <https://www.mma.gov.br/images/arquivo/80049/Areas%20Prioritarias/Caatinga/BIOMA%20CAATINGA.pdf>
- MMA (Ministério do Meio Ambiente) (2018). *Unidades de Conservação (todas)*. In Áreas Especiais. Retrieved from <https://mapas.mma.gov.br/i3geo/datadownload.htm>
- Nogueira, M. G., Oliveira, P. C. R., & Britto, Y. C. T. (2008). Zooplankton assemblages (Copepoda and Cladocera) in a cascade of reservoirs of a large tropical river (SE Brazil). *Limnetica*, 27, 151–170.
- Nolte, C., Agrawal, A., Silvius, K. M., & Soares-Filho, B. S. (2013). Governance regime and location influence avoided deforestation success of protected areas in the Brazilian Amazon. *Proceedings of the National Academy of Sciences of the United States of America*, 110(13), 4956–4961. <https://doi.org/10.1073/pnas.1214786110>
- Overbeck, G. E., Vélez-Martin, E., Scarano, F. R., Lewinsohn, T. M., Fonseca, C. R., Meyer, S. T., ... Pillar, V. D. (2015). Conservation in Brazil needs to include non-forest ecosystems. *Diversity and Distributions*, 21, 1455–1460. <https://doi.org/10.1111/ddi.12380>
- Pelicice, F. M., Azevedo-Santos, V. M., Vitule, J. R. S., Orsi, M. L., Lima-Junior, D. P., Magalhães, A. L. B., ... Agostinho, A. A. (2017). Neotropical freshwater fishes imperilled by unsustainable policies. *Fish and Fisheries*, 18, 1119–1133. <https://doi.org/10.1111/faf.12228>
- Peres, C. A. (2005). Why we need Megareserves in Amazonia. *Conservation Biology*, 19, 728–733. <https://doi.org/10.1111/j.1523-1739.2005.00691.x>
- Petere, M. Jr, Barthem, R. B., Cordoba, E. A., & Gomez, B. C. (2004). Review of the large catfish fisheries in the upper Amazon and the stock depletion of piraiba (*Brachyplatystoma filamentosum* Lichtenstein). *Reviews in Fish Biology and Fisheries*, 14, 403–414.
- Pompeu, P. S., Reis, L. S., Gandini, C. V., Souza, R. C. R., & Favero, J. M. (2009). The ichthyofauna of upper rio Capivari: Defining conservation strategies based on the composition and distribution of fish species. *Neotropical Ichthyology*, 7(4), 659–666. <https://doi.org/10.1590/S1679-62252009000400015>
- QGIS Development Team (2018). *QGIS Geographic Information System*. Open Source Geospatial Foundation Project. Retrieved from <https://www.qgis.org/en/site/>
- Saunders, D. L., Meeuwig, J. J., & Vincent, A. C. J. (2002). Freshwater protected areas: Strategies for conservation. *Conservation Biology*, 16, 30–41. <https://doi.org/10.1046/j.1523-1739.2002.99562.x>
- Simberloff, D., Martin, J.-L., Genovesi, P., Maris, V., Wardle, D. A., Aronson, J., ... Vilà, M. (2013). Impacts of biological invasions: What's what and the way forward. *Trends in Ecology & Evolution*, 28, 58–66. <https://doi.org/10.1016/j.tree.2012.07.013>
- SNUC - Sistema Nacional de Unidades de Conservação da Natureza (2000). Lei no 9.985 de 18 de Julho de 2000; decreto no 4.340 de 22 de agosto de 2002 5ª ed. aum. Brasília MMA/SBF, 2004.
- Tófoli, R. M., Dias, R. M., Alves, G. H. Z., Hoehinghaus, D. J., Gomes, L. C., Baumgartner, M. T., & Agostinho, A. A. (2017). Gold at what cost? Another megaproject threatens biodiversity in the Amazon. *Perspectives in Ecology and Conservation*, 15, 129–131. <https://doi.org/10.1016/j.pecon.2017.06.003>
- Turtle Taxonomy Working Group (2017). Turtles of the World: Annotated Checklist and Atlas of Taxonomy, Synonymy, Distribution, and Conservation Status. *Chelonian Research Monographs*, 7, 1–292.
- Vitule, J. R. S., Azevedo-Santos, V. M., Daga, V. S., Lima, D. P. Jr, Magalhães, A. L. B., Orsi, M. L., ... Agostinho, A. A. (2015). Brazil's drought: Protect biodiversity. *Science*, 347, 1427–1428. <https://doi.org/10.1126/science.347.6229.1427-b>
- Vitule, J. R. S., Freire, C. A., Vazquez, D. P., Nuñez, M. A., & Simberloff, D. (2012). Revisiting the potential conservation value of non-native species. *Conservation Biology*, 26, 1153–1155. <https://doi.org/10.1111/j.1523-1739.2012.01950.x>
- Winemiller, K. O., McIntyre, P. B., Castello, L., Fluet-Chouinard, E., Giarrizzo, T., Nam, S., ... Sáenz, L. (2016). Balancing hydropower and biodiversity in the Amazon, Congo, and Mekong. *Science*, 351, 128–129. <https://doi.org/10.1126/science.aac7082>
- Zarfl, C., Lumsden, A. E., Berlekamp, J., Tydecks, L., & Tockner, K. (2015). A global boom in hydropower dam construction. *Aquatic Sciences*, 77, 161–170. <https://doi.org/10.1007/s00027-014-0377-0>

BIOSKETCH

Author contributions: V.M.A.S. conceived the idea and wrote the first version. All authors contributed to the writing and revised all versions of the manuscript.

How to cite this article: Azevedo-Santos VM, Frederico RG, Fagundes CK, et al. Protected areas: A focus on Brazilian freshwater biodiversity. *Divers Distrib*. 2019;25:442–448. <https://doi.org/10.1111/ddi.12871>