

**The text that follows is a REPRINT  
O texto que segue é um REPRINT.**

Please cite as:  
Favor citar como:

**Laurance, W.F., S. Bergen, M.A.  
Cochrane, P.M. Fearnside, P.  
Delamonica, S. D'Angelo, C. Barber  
and T. Fernandes. 2005. The future  
of the Amazon. pp. 583-609 In: E.  
Bermingham, C.W. Dick and C.  
Moritz (eds.) *Tropical Rainforests:  
Past, Present and Future*. University  
of Chicago Press, Chicago, Illinois,  
U.S.A. 1004 pp.**

Copyright University of Chicago Press, Chicago, Illinois, U.S.A..

\  
The original publication is available from:  
A publicação original está disponível de:  
University of Chicago Press, Chicago, Illinois, U.S.A.

## The Future of the Amazon

---

WILLIAM F. LAURANCE, SCOTT BERGEN,  
MARK A. COCHRANE, PHILIP M. FEARNSIDE,  
PATRICIA DELAMÔNICA, SAMMYA AGRA D'ANGELO,  
CHRISTOPHER BARBER, AND TITO FERNANDES

### ABSTRACT

What is happening to the forests of the Amazon today? What will the region be like in another 20 years? Which areas are most vulnerable, and which stand a chance of surviving the coming decades largely intact? In this chapter we grapple with these thorny questions. If you are interested in the Amazon, but are not up to date on all the development issues, trends, and controversies, then this chapter was written for you. Parts of this chapter are speculative; in developing our predictions it was necessary to adopt a coarse-grained approach. Here we are predicting broad, basin-wide patterns of forest conversion, not specific, local-scale patterns, because such fine-scale predictions would undoubtedly prove rife with errors. We begin by summarizing the principal threats to the Amazon today: deforestation, logging, forest fragmentation, and wildfires. We then attempt to predict the pattern and scale of forest conversion over the next two decades based on current trends in deforestation, human migration, and planned and existing highways, roads, and infrastructure projects.

### INTRODUCTION

The Amazon contains nearly 60% of the world's remaining tropical rainforest (Whitmore 1997). Closed-canopy forests in the Amazon basin encompass about 5.3 million square kilometers, an area the size of western Europe (Sarre, Sobral Filho, and Reis 1996). By far the largest forest type is *terra firme*—forest that is not seasonally flooded. There also are extensive areas of seasonally flooded forest along rivers and in floodplains (termed *várzea* if they are flooded by relatively nutrient-rich white waters and *igapó* if inundated by nutrient-poor black waters), as well as limited areas of bamboo forest and vine forest. In addition, there are scattered savannas and open forests in drier areas of the basin, where narrow strips of rainforest vegetation (termed “gallery forest”) often persist along permanent rivers and streams (IBGE 1997).

Most of the Amazon is flat or undulating, occurs at low elevations (< 300 m),

and has poor soils. About four-fifths of the Amazon's soils are classified as latosols (Brown and Prance 1987; Sarre, Sobral Filho, and Reis 1996), which are typically heavily weathered, acidic, high in toxic aluminum, and poor in nutrients such as phosphorus, calcium, and potassium. In these soils, clay particles can form tight aggregations, giving the soils poor water-holding characteristics, even with high clay contents (Richter and Babbar 1991). The most productive soils in the Amazon are concentrated along the basin's western margin, in the Andean foothills and their adjoining floodplains. These areas are much more recent geologically than the rest of the basin (see Ruokolainen, Tuomisto, and Kalliola, chap. 13 in this volume) and thus their soils are far less heavily weathered.

There is a strong gradient of rainfall and seasonality in the Amazon. In general, forests in the basin's eastern and southern portions are driest, with the strongest dry season. Although evergreen, these forests are near the physiological limits of tropical rainforest and can persist only by having deep root systems that can take up groundwater during the dry season (Nepstad et al. 1994). The wettest and least seasonal forests are in the northwestern Amazon, with the central Amazon being intermediate; forests in these areas do not require deep roots.

## CURRENT THREATS TO THE AMAZON

### *Deforestation*

In the past, Amazonian development has been limited by the basin's poor soils, its remoteness from major population centers, and diseases such as malaria and yellow fever. This situation is rapidly changing. More of the Brazilian Amazon's forest has been destroyed over the last 30 years than in the previous 450 years since European colonization (Lovejoy 1999). Losses of Amazonian forests in Bolivia, Ecuador, Colombia, and Peru have also risen dramatically in recent decades (Sarre, Sobral Filho, and Reis 1996; Viña and Cavalier 1999; Steininger, Tucker, Ersts et al. 2001; Steininger, Tucker, Townshend et al. 2001).

Deforestation rates in the Amazon probably average 3–4 million ha per year—larger than the total area of Belgium. The most reliable deforestation statistics are those for the Brazilian Amazon, which have been produced annually since 1990 (except 1993) based on Landsat TM images (INPE 1996, 1998, 2000). Despite various initiatives to slow forest loss, deforestation in the Brazilian Amazon (fig. 28.1) rose significantly ( $P = .02$ , Mann-Whitney  $U$ -test) from 1990–1994 to 1995–2002 (Laurance, Albernaz, and Da Costa 2001). (Economic conditions influenced these trends; 1991 was an exceptionally low year because Brazilian bank accounts were frozen, while 1995 was an exceptionally high year be-







**Figure 28.2** Industrial logging creates labyrinths of roads that promote forest colonization and overhunting. (Photograph by W. F. Laurance.) See also plate 7.

obtained long-term forest leases (termed “concessions”: Colchester 1994; Sizer and Rice 1995). In 1996 alone, Asian companies invested more than \$500 million in the Brazilian timber industry (Muggiati and Gondim 1996). Asian multinationals now control at least 13 million ha of Amazonian forest (Laurance 1998).

A striking feature of the Amazonian timber industry is that illegal logging is rampant (see also MacKinnon, chap. 27 in this volume). A 1997 study by the Brazilian government concluded that 80% of Amazonian logging was illegal, and recent raids have netted massive stocks of stolen timber (Abramovitz 1998). Aside from widespread illegal cutting, most legal operations of the hundreds of domestic timber companies in the Amazon are poorly managed. A government inspection of thirty-four operations in Paragominas, Brazil, for example, concluded that “the results were a disaster,” and that not one was using accepted practices to limit forest damage (Walker 1996). In a controversial effort to gain better control over logging operations, Brazil opened thirty-nine of its National Forests (totaling 14 million ha) to logging in 1997, arguing that logging concessions would not be granted to companies with poor environmental records (*Critica* 1997). Much larger areas of the Brazilian Amazon are likely to be designated as logging reserves in the future (Verissimo, Cochrane, and Souza 2002).

### *Forest Fragmentation*


The rapid pace of deforestation is leading to widespread forest fragmentation. Habitat fragmentation has myriad effects on Amazonian forests, such as altering the diversity and composition of fragment biotas and changing ecological processes such as pollination, nutrient cycling, and carbon storage (Lovejoy et al. 1986; Bierregaard et al. 1992; Didham et al. 1996; Laurance and Bierregaard 1997; Laurance, Lovejoy et al. 2002). Edge effects—ecological changes associated with the abrupt, artificial edges of forest fragments—penetrate at least 300 m into Amazonian forests (Laurance et al. 1997, 1998a, 2000) and possibly much farther (Skole and Tucker 1993; Laurance 2000).

One key study found that by 1988, the area of forest in the Brazilian Amazon that was fragmented ( $< 100 \text{ km}^2$  in area) or prone to edge effects ( $< 1 \text{ km}$  from forest edge) was over 150% larger than the area that had actually been deforested (Skole and Tucker 1993). Because over 15% of the region's forest has now been cleared (INPE 1996, 1998, 2000), the total area affected by fragmentation, deforestation, and edge effects could constitute a third of the Brazilian Amazon today (Laurance 1998). This figure would probably rise if the extensive areas affected by logging and ground fires were included, but such changes are difficult to detect (Stone and Lefebvre 1998) and have not been quantified in the satellite images used to map Amazonian deforestation (Nepstad, Verissimo et al. 1999).

Forest fragmentation is occurring at many spatial scales. On a regional scale, the once remote interior of the Amazon is being dissected by major highways, powerlines, and transportation projects, which inevitably lead to rapid deforestation. On a local scale, different land uses tend to generate characteristic patterns of fragmentation. Cattle ranchers, for example, typically destroy large, rectangular blocks of forest, and habitat fragments that persist in such landscapes are somewhat regular in shape (fig. 28.3A). Forest colonization projects, however, result in more complex patterns of fragmentation, creating very irregularly shaped fragments and a high proportion of forest edge (Dale and Pearson 1997; Laurance et al. 1998b). The resulting spatial pattern has been likened to the ribs of a fish (fig. 28.3B).

### *Wildfires*

Under natural conditions, large-scale fires are very rare in Amazonian rainforests, occurring perhaps once or twice every thousand years during exceptionally severe El Niño droughts (Sanford et al. 1985; Saldariagga and West 1986; Meggers 1994; Piperno and Becker 1996). (However, some drier Amazonian for-



**Figure 28.3** Different land uses produce characteristic patterns of forest loss and fragmentation. (A) Deforestation by large-scale cattle ranching (near Paragominas, Pará). (B) “Fishbone” deforestation pattern associated with forest colonization projects (near Tailândia, Pará). Each rectangle shows an area of 570 km<sup>2</sup>. (From Cochrane and Laurance 2002.)

est formations, such as sandy-soil campinaranas, apparently burned more frequently than did primary rainforest: B. W. Nelson, personal communication.) Closed-canopy tropical forests are poorly adapted to fire (Uhl and Kauffman 1990), and even light ground fires can cause high tree mortality (Kauffman 1991; Barbosa and Fearnside 1999; Cochrane and Schulze 1999; Cochrane et al. 1999; Nepstad, Moreira, and Alencar 1999).

Fire is used commonly in the Amazon today to clear forests, destroy slash piles, and help control weeds in pastures. Over a 4-month period in 1997, satellite images revealed nearly 45,000 separate fires in the Amazon (P. Brown 1998), virtually all of them human-caused. During drought years, smoke from forest burning becomes so bad that regional airports must be closed and hospitals

report dramatic increases in the incidence of respiratory problems (Laurance 1998).

Human land uses dramatically increase the incidence of fire in tropical forests. Logged forests are far more susceptible to fires, especially during droughts. Logging increases forest desiccation and woody debris (Uhl and Kauffman 1990) and greatly increases access by slash-and-burn farmers and ranchers, which are the main sources of ignition (Uhl and Buschbacher 1985). The combination of logging, migrant farmers, and drought was responsible for the massive fires that destroyed millions of hectares of Southeast Asian forests in 1982–1983 and 1997–1998 (Leighton and Wirawan 1986; Woods 1989; N. Brown 1998; MacKinnon, chap. 27 in this volume).

Fragmented forests are also vulnerable to fire because fragment edges are prone to desiccation (Kapos 1989) and because forest remnants are juxtaposed with fire-prone pastures, farmlands, and regenerating forests. Ground fires (fig. 28.4) originating in nearby pastures can penetrate hundreds to thousands of meters into fragmented forests (Kauffman 1991; Cochrane and Laurance 2002). These low-intensity fires kill many trees and increase canopy openings and fuel loads, making the forest far more prone to catastrophic wildfires in the future (Cochrane et al. 1999; Cochrane and Schulze 1999). During the 1997–1998 El Niño drought, wildfires lit by farmers and ranchers swept through an estimated 3.4 million ha of fragmented and natural forest, savanna, regenerating forest, and farmland in the northern Amazonian state of Roraima (Barbosa and Fearnside 1999), and there were many large fires in other locations (Cochrane and Schulze 1998).

Available evidence suggests that there might be a “deforestation threshold” above which landscapes become far more prone to fires. This could potentially occur as a result of positive feedbacks among deforestation, regional drying, smoke, and fire (Cochrane and Schulze 1999; Cochrane et al. 1999; Nepstad, Moreira, and Alencar 1999; Nepstad, Verissimo et al. 1999; Laurance and Williamson 2001). Amazonian forests recycle at least half of all rainfall back into the atmosphere, helping to maintain frequent rains, lower surface temperatures, and moderate dry seasons (Salati and Vose 1984). Regional deforestation can reduce rainfall (IPCC 1996), making forests more fire-prone and, in turn, promoting additional deforestation and fires. Smoke particles from fires further reduce rainfall by trapping microdroplets of water in the atmosphere, precluding the formation of raindrops (Rosenfeld 1999). Such positive feedbacks are most likely in the drier eastern, southern, and north-central areas of the Amazon, where rainforests are already near their physiological limits (Nepstad et al. 1998).



**Figure 28.4** Low-intensity ground fires can penetrate considerable distances into forests, killing many trees and making the forests vulnerable to devastating wildfires in the future. (Photograph by M. A. Cochrane.)

### *Ancillary Threats*

Today, even the remotest areas of the Amazon are being influenced by human activities. Illegal gold mining is widespread, with wildcat miners polluting streams with mercury (used to separate gold from sediments) and degrading stream basins with pressure hoses. Illegal miners also threaten indigenous Indians through intimidation and introductions of new diseases (Christie 1997). In addition, increasing numbers of major oil, natural gas, and mineral developments are being sanctioned by Amazonian governments (Nepstad et al. 1997; Laurance 1998). Finally, hunting pressure is growing throughout the Amazon because of greater access to forests and markets and the common use of shotguns (Peres 2001). Intensive hunting can dramatically alter the structure of animal communities, extirpate species with low reproductive rates, and exacerbate the effects of habitat fragmentation on exploited species (Robinson and Redford 1991).

### FUTURE THREATS TO THE AMAZON

In this section we highlight several proximate and ultimate factors that will affect future development trends in the Amazon. These factors relate mainly to projected changes in population size, infrastructure development, and spatial patterns of forest conversion (see Laurance, Cochrane et al. 2001; Laurance, Fearnside et al. 2001).

#### *Population Growth*

The human population of the Amazon is increasing rapidly, for two reasons. First, populations have been growing throughout Latin America, nearly tripling (from 166 to 448 million) between 1950 and 1990 (Mahar and Schneider 1994). Although the traditionally high fertility rates of Latin American women have declined in recent decades, the momentum of population growth will continue for some time because a large proportion of the population is young or still in their child-bearing years. In addition, Amazonian residents often begin bearing children early—in their late teens or early twenties—which contributes substantially to rapid population growth.

Second, there is much immigration into the Amazon. In Brazil, poor economic conditions and droughts in the northeast, limited opportunities in large cities, the displacement of agricultural workers by mechanized farming, and government colonization programs designed to reduce urban overcrowding



### *New Infrastructure Projects*

Amazonian countries have ambitious short-term plans to develop major infrastructure projects encompassing large expanses of the basin. These projects are intended to accelerate economic development and exports, especially in the agriculture, timber, and mining sectors of the economy. In the Brazilian Amazon, massive investments, on the order of \$40 billion in the years 2000–2007, are being implemented to fast-track the construction of dozens of major infrastructure projects—highways, railroads, gas lines, hydroelectric projects, powerlines, and river channelization projects (Friends of the Earth 1999; *Avanço Brasil* 1999). The Amazonian road network is being rapidly expanded and upgraded, with many unpaved sections being converted to paved, all-weather highways. Key environmental agencies, such as the Ministry of the Environment, are being largely excluded from the planning of these projects (Laurance and Fearnside 1999; Laurance, Cochrane et al. 2001).

One indicator of the scale of planned development is the rapidly expanding network of hydroelectric dams. At least nineteen major (100–13,000 megawatt) dams are planned in the Brazilian Amazon over the next 10–20 years, nearly all in forested areas (Eletrobrás 1998). These new dams will vastly increase the 600,000 ha of forest that is currently inundated by reservoirs (because the region is quite flat, Amazonian hydroelectric reservoirs are often very large: Fearnside 1995). Most of these dam sites are in tributaries flowing northward into the Amazon River from Brazil's central plateau (the Tocantins, Araguaia, Xingu, and Tapajós Rivers), a region with a high concentration of indigenous peoples (Fearnside 1990). In addition to destroying forests and degrading aquatic systems, hydroelectric dams require networks of access roads and powerline clearings, which promote further forest loss and fragmentation.

New infrastructure projects will dissect vast expanses of the Amazon. The northern Amazon, for example, has already been bisected by BR-174, and will soon be cut by a road linking the rapidly growing city of Boa Vista to Guyana and by a major powerline corridor linking Guri Dam in Venezuela with Boa Vista. These projects will affect large expanses of forest as well as many indigenous groups in the northern Amazon (Soltani and Osborne 1994), and will greatly increase access to the region for loggers, ranchers, miners, and colonists. Paving of the BR-163 highway between Cuiabá and Santarém could create a large swath of degraded forest through the south-central part of Amazonia.

Other projects are equally ambitious. When completed, the massive Ferro-norte Railway will be the largest transportation project in Brazil, traversing over 4,000 km of Amazonian forest while linking the cities of Santarém (along the


Amazon River) and Porto Velho (in Rondônia) to those in southern Brazil. In the central and eastern Amazon, the Madeira, Tocantins, and Araquá rivers are being channelized in order to allow deep-water river barges to transport soybeans from rapidly expanding agricultural areas in central Brazil (Fearnside 2001). In the southern Amazon, planned road projects will traverse large expanses of forest and ascend the Andes to reach the Pacific coast, passing through Bolivia, Peru, and northern Chile. In addition, a 3,000 km natural gas line currently under construction will run from Santa Cruz, Bolivia, to São Paulo, Brazil (Soltani and Osborne 1994).

### *Logging and Mining Booms*

Increasingly, logging and mining activities are becoming important driving forces in the exploitation of the Amazonian frontier. Timber, petroleum, natural gas, and mineral resources (iron ore, bauxite, gold, copper: Sarre, Sobral Filho, and Reis 1996) provide the economic impetus for construction of roads, highways, and transportation networks, which greatly increase access to forests for colonists, ranchers, and land speculators. Roads created for oil exploration and development in Amazonian Ecuador have caused a drastic increase in forest colonization, land speculation, and commercial hunting (Holmes 1996). Similar trends are likely in the Peruvian Amazon, much of which is currently being opened for oil and gas concessions (fig. 28.5). Logging operations also greatly increase access to frontier areas; it has been estimated that 10,000 to 15,000 km<sup>2</sup> of forest are being logged each year in the Brazilian Amazon alone, a figure nearly as large as the area being deforested each year (Nepstad, Moreira, and Alencar 1999; Nepstad, Verissimo et al. 1999).

## PREDICTING THE FUTURE OF THE BRAZILIAN AMAZON

In this section we attempt to predict the scale and pattern of Amazonian forest degradation over the next two decades. We confine our predictions to the Brazilian Legal Amazon, which constitutes about two-thirds of the basin (ca. 4.9 million km<sup>2</sup>), because accurate spatial data on deforestation, transportation networks, and planned infrastructure projects for the other Amazonian and Guianan countries were very difficult to acquire on a consistent basis. Our analysis is based on a GIS (Geographic Information Systems) model that integrates spatial data on existing and planned development activities.




**Figure 28.5** Much of the Peruvian Amazon is being opened up for oil and gas exploration and development. Shaded areas show current oil and gas concessions, mostly owned by multinational corporations.

*GIS Data Layers*

To develop our model, we used the best and most recent available information on forest cover, rivers, planned and existing roads and infrastructure projects, fire-proneness of forests, logging and mining intensity, and various conservation units (table 28.1). Principal data sources for forest cover, current roads

A




-  Present paved
-  Present unpaved
-  Future unpaved
-  Future paved

SCALE 1:25724698


100 0 100 200



KILOMETERS



B



-  Present utilities
-  Future utilities
-  Present railroads
-  Future railroads
-  Future channels
-  Present dams
-  Future dams

**Figure 28.6** Existing and planned highways and infrastructure projects in the Brazilian Amazon. (A) Highways and roads. (B) Major infrastructure projects. “Utilities” are gas lines and powerlines, while “channels” are river channelization projects.

utility. The probability of forest fires was based on the map of Nepstad et al. (1998; Nepstad, Moreira, and Alencar 1999), who integrated extensive data on forest cover, seasonal soil water availability, recent fires, and logging activity in order to predict areas of high, moderate, and low fire vulnerability during the 1998 dry season. Maps of the estimated extent of logging (both legal and illegal), industrial mining, and illegal gold mining were produced by IBAMA (Brazil's national environmental agency) in 1998.

The Brazilian Amazon has a variety of federal and state conservation units that vary considerably in their degree of environmental protection. We identified thirteen major types of reserves and parks, which we placed into three general categories (table 28.2). Areas with nominally high protection include National Parks, Ecological Stations, and Ecological Reserves, which nominally receive strong protection. Sanctioned activities in such areas include research, education, and often, recreation and tourism.

Areas with moderate protection are National Forests, Extractive Reserves, and Sustainable Development Reserves, among others, which may be legally subjected to nominally "sustainable" levels of industrial logging, agriculture, livestock grazing, hunting, fishing, tourism, and extraction of non-timber products (e.g., rubber, fuelwood, fruits, seeds, fibers). Mining is usually prohibited, although illegal gold mining certainly occurs in some of these reserves.

The final category, areas with uncertain protection, includes the extensive indigenous lands and reserves that collectively constitute about 18% of the Brazilian Legal Amazon. In some areas, these lands may be more effectively protected than National Parks, especially where indigenous people are territorial and repel illegal colonists, loggers, and gold miners. In other areas, however—particularly where indigenous people have frequent contact with outsiders—a corruption of traditional lifestyles can occur. In a number of cases, indigenous groups in Brazil have sold their timber to commercial loggers, permitted wild-cat mining, overhunted wildlife, illegally cleared protected lands, invaded national parks, impeded firefighters, and even assaulted government inspectors attempting to control illegal logging (Redford and Stearman 1993; Alvard et al. 1997; Margolis 2000). Hence, environmental protection in indigenous lands and reserves is likely to be highly variable, and will tend to decline as contact with outsiders increases.

### *Buffer Zones*

Roads and infrastructure projects promote forest degradation by greatly increasing human access, and in some cases (such as hydroelectric reservoirs) by

**Table 28.2** Legally permitted activities within conservation areas in the Brazilian Amazon

Type of area	Recreation & tourism	Agriculture & livestock	Logging	Non-timber harvests	Hunting	Mining
<b>Areas with nominally high protection</b>						
National/state parks	Yes	No	No	No	No	No
Ecological reserves	Yes	No	No	No	No	No
Biological reserves	No	No	No	No	No	No
Ecological stations	No	No	No	No	No	No
<b>Areas with moderate protection</b>						
National/state forests	Yes	Yes	Yes	Yes	Yes <sup>a</sup>	No
National forest reserves	Yes	Yes	Yes	Yes	Yes <sup>a</sup>	No
Extractive reserves	Yes	Yes	Yes	Yes	Yes <sup>a</sup>	No
State extractive forests	Yes	Yes	Yes	Yes	Yes <sup>a</sup>	No
Sustainable use forests	Yes	Yes	Yes	Yes	Yes <sup>a</sup>	No
Sustainable development reserves	Yes	Yes <sup>a</sup>	Yes	Yes	Yes <sup>a</sup>	No
Environmental protection areas	Yes	Yes <sup>b</sup>	Yes <sup>b</sup>	Yes <sup>b</sup>	No	Yes <sup>b</sup>
Areas of relevant ecological interest	Yes	Yes <sup>b</sup>	No	Yes <sup>b</sup>	No	No
<b>Areas with uncertain protection</b>						
Indigenous lands and reserves	No	Yes	Yes	Yes	Yes	No

*Sources:* Silva 1996; Olmos, Queiroz Filho, and Lisboa 1998; Rylands and Pinto 1998; Borges et al. 2001; Web sites of IBAMA, Instituto Socioambiental, and IBGE; personal communication with Luciene Pohl of Brazil's National Indian Foundation (FUNAI).

<sup>a</sup>Hunting is allowed in some areas; for others information was unavailable.

<sup>b</sup>These activities are not expressly permitted, but because people are allowed to live in these reserves they will certainly occur, at least on a limited scale.

destroying large areas of forest directly. To predict the future effects of planned roads and other infrastructure projects, we assessed the past effects of existing highways and roads on primary forest cover in the Amazon.


To do this, we overlaid the existing road network on the Landsat TM-based Pathfinder map of the Brazilian Amazon for 1992. Many of the region's major highways (e.g., Belém-Brasília, Transamazon, BR-364) were constructed in the 1960s and 1970s, and thus had been in existence for 15–25 years by 1992—roughly comparable to the 20-year time frame for our predictions. Initially, five “buffer zones” were created around all paved highways (0–10, 11–25, 26–50, 51–75, and 76–100 km on each side of the highway), and the percentage loss of primary forest cover within each zone was determined. This analysis was then repeated using the entire network of highways and unpaved roads. Clouds, cloud shadows, and rivers were removed from the analyses (< 5% of total area). Buffers were truncated if they passed outside the Brazilian Legal Amazon. Deforestation was registered only for closed-canopy forests; losses of other habitats (e.g., savanna) were not included. Analyses were run on a Silicon Graphics Origin supercomputer at the Basic Science and Remote Sensing Initiative, Michigan State University.

As expected, the analyses (fig. 28.7) revealed that deforestation strongly increased near highways and roads. Both averaged about 30% forest loss within the 0–10 km buffer zone, but highways had more far-reaching effects than roads, averaging about 20% and 15% forest loss in the 11–25 and 26–50 km zones, respectively. Roads tended to generate more localized deforestation, with average forest loss declining below 15% beyond 25 km from the road.

Networks of roads tend to proliferate near highways, as is evident, for example, along the Belém-Brasília and eastern Transamazon highways. The most far-reaching effect we observed was the construction of 200–300 km long state and local roads ramifying laterally from highways in Pará, Mato Grosso, and Amazonas states (see fig. 28.6). Road networks are also generated by infrastructure projects, as it is nearly impossible to construct hydroelectric dams, powerlines, gas lines, and other major facilities without road access.

### *Model Assumptions*

We used the buffer zone analyses to help generate two alternative predictions—termed the “optimistic” and “non-optimistic” scenarios—for the future of the Brazilian Amazon. Our models predict the spatial distribution of four broad land use categories. The first category is heavily degraded areas, regions in which primary forest cover is likely to be absent or markedly reduced and



**Figure 28.7** Percentage of closed-canopy forest destroyed by 1992 as a function of distance from paved highways, and from all roads and highways, in the Brazilian Amazon.

heavily fragmented. Such areas are highly vulnerable to edge effects, fires, logging, and overhunting and are severely degraded ecologically. Moderately degraded areas have mostly intact primary forest cover (> 85%), but contain localized forest clearings and some roads and may be affected by logging, mining, hunting, and oil and gas exploration. Lightly degraded areas have nearly intact primary forest cover (> 95%), but can experience illegal gold mining, small-scale farming, hunting, hand logging, and non-timber resource extraction (e.g., rubber tapping). “Pristine areas” have fully intact primary forest cover and are free from anthropogenic impacts aside from limited hunting, fishing, and swidden farming by traditional indigenous communities.

The optimistic and non-optimistic models differ in that the former assumes that highways, roads, and infrastructure projects will generate more localized

**Table 28.3** Explicit assumptions of "optimistic" and "non-optimistic" GIS models used to predict forest degradation in the Brazilian Amazon

	Optimistic scenario	Non-optimistic scenario
<b>Buffers around paved highways (current and planned)</b>		
Heavily degraded zone	0–25 km	0–50 km
Moderately degraded zone	26–50 km	51–100 km
Lightly degraded zone	51–75 km	101–200 km
Pristine zone	> 75 km	> 200 km
<b>Buffers around unpaved roads, railroads, powerlines, gas lines, industrial mines, and river channelization projects (current and planned)</b>		
Heavily degraded zone	0–10 km	0–25 km
Moderately degraded zone	11–25 km	26–50 km
Lightly degraded zone	26–50 km	51–100 km
Pristine zone	> 50 km	> 100 km
<b>Buffers around hydroelectric reservoirs</b>		
Heavily degraded zone	Area inundated	Area inundated
Moderately degraded zone	0–5 km	0–10 km
Lightly degraded zone	6–10 km	11–25 km
Pristine zone	> 10 km	> 25 km
Areas prone to logging	Moderately degraded	Moderately degraded
Areas prone to wildcat mining	Lightly degraded	Lightly degraded
Areas prone to fires		
High vulnerability	Moderately degraded	Heavily degraded
Moderate vulnerability	Lightly degraded	Moderately degraded
<b>Conservation areas</b>		
High-protection areas outside buffers	Pristine	Pristine
High-protection areas inside buffers	Pristine	Lightly degraded
Moderate-protection areas outside buffers	Lightly degraded	Lightly degraded
Moderate-protection areas inside buffers	Lightly degraded	Moderately degraded
Indigenous areas outside buffers	Pristine	Lightly degraded
Indigenous areas inside buffers	Lightly degraded	Moderately degraded

effects and that conservation areas will be less prone to disturbances (table 28.3). The sizes of the buffer zones used in the models were necessarily somewhat arbitrary, but have an empirical basis in our analyses of past deforestation. In the non-optimistic model, for example, we assumed that paved highways would create a 50 km wide zone of heavily degraded forest on each side because our analysis suggested that these areas averaged less than 85% forest cover (see fig. 28.7). Such areas would be prone to logging, fragmentation, fires, edge effects, and other ecological changes that could affect much of the remaining forest cover (see Skole and Tucker 1993; Nepstad, Verissimo et al. 1999, Cochrane et al. 1999; Gascon, Williamson, and da Fonseca 2000; Laurance 2000). Likewise, we conservatively assumed that the lightly degraded zone would extend 100–200 km from paved highways because we observed a number of roads stretching at least 200 km from existing highways.

In both scenarios, logging and wildcat mining were assumed to cause moderate and light forest degradation, respectively. However, the models differed in terms of the viability of protected areas. The optimistic model assumed that all reserves would remain pristine or only lightly degraded, whereas the non-optimistic model assumed that indigenous lands and reserves with moderate protection (or parts thereof) would be moderately degraded within 50 km of roads or 100 km of highways; otherwise they would be pristine or lightly degraded. The non-optimistic model also assumed that reserves with high protection would be lightly degraded near roads and highways.

### *GIS Analyses*


All maps and spatial data were georeferenced to a geographic coordinate system, using Imagine 8.3 software (ERDAS 1998). Subsequently, georeferenced digital images were used for vector data layer construction, using Arc/Info 7.2.1 (ESRI 1999) via heads-up digitizing methods. Road and infrastructure buffers were created with Arc/Info software for the appropriate distances. Data layers were integrated with overlay methodology. Most analyses were performed on a Silicon Graphics Indigo2 workstation at Biological Dynamics of Forest Fragments Project (BDFFP) headquarters in Manaus, Brazil.

### *Results*


The optimistic scenario (fig. 28.8A) suggests that by the year 2020 there will be continued deforestation in the southern and eastern portions of the Brazilian Amazon and considerable large-scale fragmentation of forests in the central and southern parts of the basin. The Brazilian Amazon will be nearly bisected by heavily to moderately degraded areas along a north-south axis running from Rondônia to Manaus and northward to Venezuela. Pristine and lightly degraded forests will be fragmented into several blocks, with the largest tract surviving in the western Brazilian Amazon. According to this scenario, pristine forests will constitute just 28.0% of the region, and lightly degraded forests will constitute another 28.0%. Almost 27% of the region will be deforested or heavily degraded (fig. 28.9).

**Figure 28.8** Projected forest degradation in the Brazilian Amazon by the year 2020: (A) optimistic scenario; (B) non-optimistic scenario. Gray areas = deforested or heavily degraded, including savannas and other nonforested areas; crosshatched areas = moderately degraded; single-hatched areas = lightly degraded; white areas = pristine.


A




B



*With new highways and infrastructure*



*Without new highways and infrastructure*



The non-optimistic scenario (fig. 28.8B) projects an even more dramatic loss of forests along the southern and eastern areas of the basin. Large-scale fragmentation is also more extensive, with much forest in the central, northern, and southeastern areas persisting only in isolated tracts. The basin is almost completely bisected by a swath of heavily degraded lands along the north-south axis running from Rondônia to Venezuela. There are very few areas of pristine forest aside from those in the western quarter of the region. This scenario predicts that pristine forests will constitute just 4.9% of the region, with lightly degraded forests constituting another 25.4%. Over 40% of the region will be heavily degraded.

### *Discussion and Implications*

Both of our models suggest that the Brazilian Amazon will be drastically altered by current development plans and prevailing land use trends over the next 20 years. The principal differences between the models are in the extent of forest loss and fragmentation (see fig. 28.8) and the relative proportions of heavily degraded versus pristine forests (see fig. 28.9).

Some degree of oversimplification in our models was inevitable. For example, we did not incorporate the effect of population density into our models, in part because we observed that local road density in the Amazon seemed to be a reasonably good surrogate for local population density. It is also apparent that the buffer zones around roads, highways, and infrastructure projects will be more variable spatially than is indicated in our models. While we have incorporated many of the factors that are likely to influence local deforestation (e.g., distance to roads, road quality [paved vs. unpaved], presence and type of protected areas, vulnerability to forest fires, logging and mining activity), it is impossible to include every potentially relevant factor (see Laurance, Albernaz et al. 2002 for further analyses of factors that influence Amazonian deforestation).

The optimistic and non-optimistic scenarios vary considerably, and it is therefore important to ask which is the most realistic. At least two considerations suggest that the non-optimistic scenario may better approximate reality. First, the non-optimistic model realistically assumes that forests with high fire vulnerability will become heavily degraded, while those of moderate vulnerability will become moderately degraded. The model of fire vulnerability we used

**Figure 28.9** Predicted percentages of Brazilian Amazon forest in four degradation classes by the year 2020 according to the optimistic and non-optimistic scenarios.

(Nepstad et al. 1998) was produced for a normal dry season and is therefore conservative, in the sense that much larger areas of the Amazon will become vulnerable to fires during periodic El Niño droughts (Nepstad et al. 1998; Cochrane and Schulze 1998, 1999).

Second, the non-optimistic model assumes that protected areas within 50 km of highways and roads will be lightly to moderately degraded. In fact, many protected areas in the Amazon are little more than "paper parks" with inadequate protection. Ferreira et al. (1999) evaluated eighty-six federal parks and protected areas in Brazil and found that 43% were at high to extreme risk because of illegal deforestation, colonization, hunting, isolation of the reserve from other forest areas, and additional forms of encroachment. More than half of all reserves (54.6%) were judged to have nearly nonexistent management. For some reserves, even our non-optimistic scenario may be overly optimistic.

The fates of indigenous lands and reserves will have an important effect on forest conservation. In many of these areas, the quality of environmental protection has declined markedly as indigenous groups come into more frequent contact with outsiders and traditional lifestyles are lost (Margolis 2000). In this sense, reducing construction of new roads and highways near indigenous lands may be one of the most effective measures to ensure that traditional management systems are not corrupted.

Other investigators have also attempted to predict spatial patterns of Amazonian deforestation. Recent studies by a Brazilian nongovernmental organization attempted to predict the extent of deforestation that will be caused by new highway construction under the *Avança Brasil* program (Nepstad et al. 2000, 2001; Carvalho et al. 2001), but did not consider the effects of other infrastructure projects (hydroelectric reservoirs, powerlines, gas lines, railroads, river channelization projects, and their associated road networks) on forests (see Fearnside 2002). In addition, an ongoing study by C. Souza Jr. (personal communication) is using data on existing road networks, logging, and recent fires in order to assess conversion pressure on Brazilian Amazonian forests. Earlier studies, such as those of Kangas (1990) and Bryant, Nielsen, and Tanglely (1997), did not incorporate the effects of massive planned highway and infrastructure developments under *Avança Brasil*, and hence are seriously out of date.

Obviously, our models illustrate only two of a potentially infinite number of possible futures for the Brazilian Amazon. While we believe our approach is based on realistic assumptions, it has two limitations. First, our model predictions are somewhat difficult to test and verify, especially for lightly degraded forests. Low-intensity selective logging and illegal gold mining, for example, are nearly impossible to detect with remote sensing, although technological im-

provements could change this in the future. Second, our models rely on specific assumptions about the future drivers of forest degradation (see table 28.3). Perhaps the most crucial assumption is that current infrastructure projects will proceed as planned and that there will be no major new development initiatives. The unforeseen construction of a new highway, for example, could alter the scale and spatial pattern of forest degradation, reducing the accuracy of our predictions.

In the Amazon, hundreds of millions of dollars are currently being expended on efforts to promote conservation planning via international programs such as the Pilot Program to Protect the Brazilian Rainforest, bilateral initiatives, and the activities of nongovernmental organizations (Laurance and Fearnside 1999; Laurance, Cochrane et al. 2001). The most important conclusion of our study is that current domestic and international efforts to promote conservation planning in the Brazilian Amazon are likely to be swamped by short-term plans to invest over US\$40 billion in Amazonian transportation and infrastructure projects. The environmental impacts of these projects will be further magnified by population increases, forest colonization projects, and rapidly expanding logging and mining industries. If our models provide even a rough approximation of the future, the forests of the Amazon will be profoundly altered over the next two decades. We conclude that a fundamental reevaluation of the criteria used in selecting, planning, and licensing large-scale development projects is urgently needed (see Fearnside 2002). Without major policy changes, current development schemes are likely to have dire effects on Amazonian forests.

#### ACKNOWLEDGMENTS

We thank Bruce Nelson, Heraldo Vasconcelos, and Rita Mesquita for discussion and reviewing the manuscript. Support was provided by the NASA-LBA program, Andrew W. Mellon Foundation, World Wildlife Fund-U.S., MacArthur Foundation, National Institute for Amazonian Research, and the Smithsonian Institution. This chapter is publication number 316 in the BDFFP technical series.

## LITERATURE CITED

- Abramovitz, J. 1998. Taking a stand: cultivating a new relationship with the world's forests. World Watch Institute, Washington, D.C.
- Alvard, M. S., J. G. Robinson, K. H. Redford, and H. Kapland. 1997. The sustainability of subsistence hunting in the neotropics. *Conservation Biology* **11**:977-982.
- Anon. 1997. Controle sobre florestas exige a reforma do IBAMA. *A Crítica Newspaper*, Manaus, Brazil, 19 January 1997.
- Anon. 1999. O debate sobre políticas públicas para Amazônia na imprensa brasileira, Vol. IV. Friends of the Earth, Amazonia Program, Brasília, Brazil.
- Avança Brasil. 1999. Avança Brasil: development structures for investment. Ministry for Development, Industry, and Foreign Trade, Brasília, Brazil.
- Barbosa, R. I., and P.M. Fearnside. 1999. Incêndios na Amazônia brasileira: estimativa da emissão de gases do efeito estufa pela queima de diferentes ecossistemas de Roraima na passagem do evento "El Niño" (1997/98). *Acta Amazonica* **29**:513-534.
- Bennett, E. L., and M. T. Gumal. 2001. The interrelationships of commercial logging, hunting, and wildlife in Sarawak: recommendations for forest management. Pages 359-274 in R. Fimbel, A. Grajal, and J. G. Robinson, editors. *The cutting edge: conserving wildlife in logged tropical forests*. Columbia University Press, New York.
- Bierregaard, R. O., Jr., T. E. Lovejoy, V. Kapos, A. dos Santos, and R. W. Hutchings. 1992. The biological dynamics of tropical rainforest fragments. *Bioscience* **42**:859-866.
- Borges, S. H., M. Pinheiro, A. Murchie, and C. Durigan. 2001. Preservação do Rio Negro: As unidades de conservação. Pages 303-329 in A. Oliveira and D. Daly, editors. *As florestas do Rio Negro*. Universidade Paulista Press, São Paulo, Brazil.
- Brown, K. S. 1987. Soils and vegetation. Pages 19-45 in T. C. Whitmore and G. T. Prance, editors. *Biogeography and Quaternary history in tropical America*. Oxford Monographs in Biogeography 3, Oxford, England.
- Brown, N. 1998. Out of control: fires and forestry in Indonesia. *Trends in Ecology and Evolution* **13**:41.
- Brown, P. 1998. Forest fires: setting the world ablaze. *The Guardian*, London, England, 20 March 1998.
- Bryant, D., D. Nielsen, and L. Tanglely. 1997. The last frontier forests: ecosystems

- and economies on the edge. World Resources Institute, Washington, D.C.
- Carvalho, G., A. C. Barros, P. Moutinho, and D. C. Nepstad. 2001. Sensitive development could protect the Amazon instead of destroying it. *Nature* **409**:131.
- Cassia, de, R. 1997. BR-174: FHC anuncia abertura de nova fronteira agrícola no Norte. *Amazonas em Tempo*, Manaus, Brazil, p. A-4, 25 June 1997.
- Christie, M. 1997. Yanomami Indians appeal for help against invaders. Reuters News Service, 31 August 1997.
- Cochrane, M. A., A. Alencar, M. D. Schulze, C. M. Souza, D. C. Nepstad, P. Lefebvre, and E. Davidson. 1999. Positive feedbacks in the fire dynamics of closed canopy tropical forests. *Science* **284**:1832-1835.
- Cochrane, M. A., and W. F. Laurance. 2002. Fire as a large-scale edge effect in Amazonian forests. *Journal of Tropical Ecology* **18**:311-325.
- Cochrane, M. A., and M. D. Schulze. 1998. Forest fires in the Brazilian Amazon. *Conservation Biology* **12**:948-950.
- Cochrane, M. A., and M. D. Schulze. 1999. Fire as a recurrent event in tropical forests of the eastern Amazon: effects on forest structure, biomass, and species composition. *Biotropica* **31**:2-16.
- Colchester, M. 1994. The new sultans: Asian loggers move in on Guyana's forests. *The Ecologist* **24**:45-52.
- Dale, V. H., and Pearson, S. M. 1997. Quantifying habitat fragmentation due to land-use change in Amazonia. Pages 400-409 *in* W. F. Laurance and R. O. Bierregaard, editors. *Tropical forest remnants: ecology, management, and conservation of fragmented communities*. University of Chicago Press, Chicago, Illinois, USA.
- Didham, R. K., J. Ghazoul, N. E. Stork, and A. J. Davis. 1996. Insects in fragmented forests: a functional approach. *Trends in Ecology and Evolution* **11**:255-260.
- Eletrobrás. 1998. Eletrobrás: The ten-year expansion plan, 1998-2007. Centrais Elétricas do Brasil, Rio de Janeiro, Brazil.
- ERDAS. 1998. Environmental research data analysis system. ERDAS, Atlanta, Georgia, USA.
- ESRI. 1999. Arc/Info 7.2.1. Environmental Systems Research Institute, Redlands, California, USA.
- Fearnside, P. M. 1987. Causes of deforestation in the Brazilian Amazon. Pages 37-61 *in* R. F. Dickinson, editor. *The geophysics of Amazonia: vegetation*

and climate interactions. John Wiley, San Francisco, USA.

- Fearnside, P. M. 1990. Environmental destruction in the Amazon. Pages 179-225 *in* D. Goodman and A. Hall, editors. The future of Amazonia: destruction or sustainable development? MacMillan, London, England.
- Fearnside, P. M. 1993. Deforestation in the Brazilian Amazon: the effect of population and land tenure. *Ambio* **8**:537-545.
- Fearnside, P. M. 1995. Hydroelectric dams in the Brazilian Amazon as sources of "greenhouse" gases. *Environmental Conservation* **22**:7-19
- Fearnside, P. M. 1997. Protection of mahogany: a catalytic species in the destruction of rain forests in the American tropics. *Environmental Conservation* **24**:303-306.
- Fearnside, P. M. 1999. Biodiversity as an environmental service in Brazil's Amazonian forests: risks, value and conservation. *Environmental Conservation* **26**:305-321.
- Fearnside, P. M. 2001. Soybean cultivation as a threat to the environment in Brazil. *Environmental Conservation* **28**:23-38.
- Fearnside, P. M. 2002. Avança Brasil: environmental and social consequences of Brazil's planned infrastructure in Amazonia. *Environmental Management* **30**:735-747.
- Ferreira, L. V., R. M. L. de Sá, R. Buschbacher, G. Batmanian, B. R. Bensuran, and K. L. Costa. 1999. Áreas protegidas ou espaços ameaçados? World Wide Fund for Nature, Brasília, Brazil.
- Gascon, C., G. B. Williamson, and G. A. B. da Fonseca. 2000. Receding forest edges and vanishing reserves. *Science* **288**:1356-1358.
- Holmes, B. 1996. The low-impact road. *New Scientist* **151(2048)**:43.
- IBGE. 1997. Diagnóstico ambiental da Amazônia Legal. CD-ROM with GIS version of RADAM maps. Brazilian Institute for Geography and Statistics (IBGE), Brasília, Brazil.
- INPE. 1996. Deforestation estimates in the Brazilian Amazon, 1992-1994. National Institute for Space Research (INPE), São Jose dos Campos, Brazil.
- INPE. 1998. Deforestation estimates in the Brazilian Amazon, 1995-1997. National Institute for Space Research (INPE), São Jose dos Campos, Brazil.
- INPE. 2000. Deforestation estimates in the Brazilian Amazon, 1998-1999. National Institute for Space Research (INPE), São Jose dos Campos, Brazil.
- IPCC. 1996. Climate change 1995: impacts, adaptations and mitigation of climate

change: scientific technical analyses. Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, England.

- Johns, A. 1997. Timber production and biodiversity conservation in tropical rain Forests. Cambridge University Press, Cambridge, England.
- Kapos, V. 1989. Effects of isolation on the water status of forest patches in the Brazilian Amazon. *Journal of Tropical Ecology* **5**:173-185.
- Kangas, P. 1990. Deforestation and diversity of life zones in the Brazilian Amazon: a map analysis. *Ecological Modeling* **49**:267-275.
- Kauffman, J. B. 1991. Survival by sprouting following fire in tropical forests of the eastern Amazon. *Biotropica* **23**:219-224.
- Laurance, W. F. 1998. A crisis in the making: responses of Amazonian forests to land use and climate change. *Trends in Ecology and Evolution* **13**:411-415.
- Laurance, W. F. 2000. Do edge effects occur over large spatial scales? *Trends in Ecology and Evolution* **15**:134-135.
- Laurance, W. F. 2001. Tropical logging and human invasions. *Conservation Biology* **15**:4-5.
- Laurance, W. F., and R. O. Bierregaard, Jr., editors. 1997. Tropical forest remnants: ecology, management and conservation of fragmented communities. University of Chicago Press, Chicago, Illinois, USA.
- Laurance, W. F., A. Albernaz, and C. Da Costa. 2001a. Is deforestation accelerating in the Brazilian Amazon? *Environmental Conservation* **28**:305-311.
- Laurance, W. F., A. K. M. Albernaz, G. Schroth, P. M. Fearnside, E. Ventincinque, and C. Da Costa. 2002. Predictors of deforestation in the Brazilian Amazon. *Journal of Biogeography* **29**:737-748.
- Laurance, W. F., M. A. Cochrane, S. Bergen, P. M. Fearnside, P. Delamonica, C. Barber, S. D'Angelo, and T. Fernandes. 2001b. The future of the Brazilian Amazon. *Science* **291**:438-439.
- Laurance, W. F., P. Delamonica, S. G. Laurance, H. Vasconcelos, and T. E. Lovejoy. 2000. Rainforest fragmentation kills big trees. *Nature* **404**:836.
- Laurance, W. F., and P. M. Fearnside. 1999. Amazon burning. *Trends in Ecology and Evolution* **14**:457.
- Laurance, W. F., P. M. Fearnside, M. A. Cochrane, S. D'Angelo, S. Bergen, and P. Delamonica. 2001c. Development of the Brazilian Amazon. *Science* **292**:1652-1654.

- Laurance, W. F., L. V. Ferreira, J. M. Rankin-de Merona, and S. G. Laurance. 1998a. Rain forest fragmentation and the dynamics of Amazonian tree communities. *Ecology* **79**: 2032-2040.
- Laurance, W. F., S. G. Laurance, and P. Delamonica. 1998b. Tropical forest fragmentation and greenhouse gas emissions. *Forest Ecology and Management* **110**:173-180.
- Laurance, W. F., S. G. Laurance, L. V. Ferreira, J. Rankin-de Merona, C. Gascon, and T. E. Lovejoy. 1997. Biomass collapse in Amazonian forest fragments. *Science* **278**: 1117-1118.
- Laurance, W. F., T. E. Lovejoy, H. L. Vasconcelos, E. M. Bruna, R. K. Didham, P. C. Stouffer, C. Gascon, R. O. Bierregaard, S. G. Laurance, and E. Sampaio. 2002. Ecosystem decay of Amazonian forest fragments: a 22-year investigation. *Conservation Biology* **16**:605-618.
- Laurance, W. F., and G. B. Williamson. 2001. Positive feedbacks among forest fragmentation, drought, and climate change in the Amazon. *Conservation Biology* **15**:1529-1535.
- Leighton, M. 1986. Catastrophic drought and fire in Borneo tropical rain forest associated with the 1982-1983 El Niño Southern Oscillation Event. Pages 75-102 *in* G. T. Prance, editor. *Tropical rain forests and the world atmosphere*. Cambridge University Press, Cambridge, England.
- Lovejoy, T. E. 1999. Preface. *Biological Conservation* **91**:100.
- Lovejoy, T. E., R. O. Bierregaard, A. Rylands, J. R. Malcolm, C. Quintela, L. Harper, K. Brown, A. Powell, G. Powell, H. Schubart, and M. Hays. 1986. Edge and other effects of isolation on Amazon forest fragments. Pages 257-285 *in* M. E. Soule, editor. *Conservation biology: the science of scarcity and diversity*. Sinauer, Sunderland, Massachusetts, USA.
- Mahar, D., and R. Schneider. 1994. Incentives for tropical deforestation: some examples from Latin America. Pages 159-171 *in* K. Brown and D. W. Pearce, editors. *The causes of tropical deforestation: the economic and statistical analysis of factors giving rise to the loss of tropical forests*. University College London Press, London, England.
- Margolis, M. 2000. Not as green as they seem. *Newsweek International*, 27 March, pp. 10-14.
- Meggers, B. J. 1994. Archeological evidence for the impact of mega-Niño events on Amazonian during the past two millennia. *Climatic Change* **28**:321-338.
- Muggiati, A. and A. Gondim. 1996. *Madeireiras*. O Estado de S. Paulo, São Paulo, Brazil, 16 September 1996.
- Nepstad, D. C., J. P. Capobianco, A. C. Barros, G. Carvalho, P. Moutinho, P.

- Lefebvre, and U. Lopes. 2000. *Avança Brasil: cenários futuros para a Amazônia*. Institute for Environmental Research in the Amazon (IPAM), Belém, Brazil.
- Nepstad, D. C., C. Carvalho, E. Davidson, P. Jipp, P. Lefebvre, G. Negreiros, E. Silva, T. Stone, S. Trumbore, and S. Vieira. 1994. The role of deep roots in the hydrological cycles of Amazonian forests and pastures. *Nature* **372**:666-669.
- Nepstad, D., G. Carvalho, A. Barros, A. Alencar, J. Capobianco, J. Bishop, P. Moutinho, P. Lefebvre, and U. Silva, Jr. 2001. Road paving, fire regime feedbacks, and the future of Amazon forests. *Forest Ecology and Management* **154**:295-407.
- Nepstad, D. C., C. Klink, C. Uhl, I. Viera, P. LeFebvre, M. Pedlowski, E. Matricardi, G. Negreiros, I. Brown, E. Amaral, A. Homma, and R. Walker. 1997. Land-use in Amazonia and the cerrado of Brazil. *Ciencia e Cultura* **49**:73-86.
- Nepstad, D. C., A. G. Moreira, and A. A. Alencar. 1999a. Flames in the rain forest: origins, impacts, and alternatives to Amazonian fires. Pilot Program to Conserve the Brazilian Rain Forest, Brasília, Brazil.
- Nepstad, D. C., A. Moreira, A. Verissimo, P. Lefebvre, P. Schlesinger, C. Potter, C. Nobre, A. Setzer, T. Krug, A. Barros, A. Alencar, and J. Pereira. 1998. Forest fire prediction and prevention in the Brazilian Amazon. *Conservation Biology* **12**:951-955.
- Nepstad, D. C., A. Verissimo, A. Alencar, C. Nobre, E. Lima, P. Lefebvre, P. Schlesinger, C. Potter, P. Moutinho, E. Mendoza, M. Cochrane, and V. Brooks. 1999b. Large-scale impoverishment of Amazonian forests by logging and fire. *Nature* **398**:505-508.
- Olmos, F., A. P. Queiroz-Filho, and C. A. Lisboa. 1998. *As unidades de conservação de Rondônia*. Secretariate for Planning (SEPLAN), Rondônia, Brazil.
- Peres, C. A. 2001. Synergistic effects of subsistence hunting and habitat fragmentation on Amazonian forest vertebrates. *Conservation Biology* **15**:1490-1505.
- Piperno, D. R., and P. Becker. 1996. Vegetational history of a site in the central Amazon Basin derived from phytolith and charcoal records from natural soils. *Quaternary Research* **45**:202-209.
- Redford, K. H., and A. M. Stearman. 1993. Forest-dwelling native Amazonians and the conservation of biodiversity. *Conservation Biology* **7**:248-255.
- Richter, D. D., and L. I. Babbar. 1991. Soil diversity in the tropics. *Advances in Ecological Research* **21**:315-389.
- Robinson, J., and K. Redford, editors. 1991. *Neotropical wildlife use and conservation*. University of Chicago Press, Chicago, Illinois, USA.

- Rosenfeld, D. 1999. TRMM observed first direct evidence of smoke from forest fires inhibiting rainfall. *Geophysical Research Letters* **26**:3105-3108.
- Rylands, A. B. 1991. The status of conservation areas in the Brazilian Amazon. World Wildlife Fund, Washington, D.C.
- Rylands, A. B., and L. P. Pinto. 1998. Conservação da biodiversidade na Amazônia Brasileira: uma análise do sistema de unidades de conservação. Brazilian Foundation for Sustainable Development, Brasília, Brazil.
- Salati, E. and P. B. Vose. 1984. Amazon basin: a system in equilibrium. *Science* **225**:129-138.
- Saldariagga, J., and D. C. West. 1986. Holocene fires in the northern Amazon basin. *Quaternary Research* **26**:358-366.
- Sanford, R. L., J. Saldariagga, K. Clark, C. Uhl, and R. Herrera. 1985. Amazon rain-forest fires. *Science* **227**:53-55.
- Sarre, A., M. Sobral Filho, and M. Reis. 1996. The amazing Amazon. ITTO Tropical Forest Update **6(4)**:3-7.
- Silva, L. L. 1996. Ecologia: manejo de áreas silvestres. Ministry of the Environment, Brasília, Brazil.
- Sizer, N., and R. Rice. 1995. Backs to the wall in Suriname: forest policy in a country in crisis. World Resources Institute, Washington, D.C., USA.
- Skole, D., and C. J. Tucker. 1993. Tropical deforestation and habitat fragmentation in the Amazon: satellite data from 1978 to 1988. *Science* **260**:1905-1910.
- Soltani, A., and T. Osborne. 1994. Arteries for global trade, consequences for Amazonia. Amazon Watch, Malibu, California, USA.
- Steininger, M. K., C. Tucker, P. Ersts, T. Killeen, and S. Hecht. 2001a. Clearance and fragmentation of semi-deciduous tropical forest in the Tierras Bajas zone, Santa Cruz, Bolivia. *Conservation Biology* **15**:856-866.
- Steininger, M. K., C. Tucker, J. Townshend, V. Bell, A. Desch, P. Ersts, and T. Killeen. 2001b. Tropical Deforestation in the Bolivian Amazon. *Environmental Conservation* **28**:127-134.
- Stone, T., and P. Lefebvre. 1998. Using multi-temporal satellite data to evaluate selective logging in Pará, Brazil. *International Journal of Remote Sensing* **19**:2517-2526.
- Uhl, C., P. Barreto, A. Verissimo, E. Vidal, P. Amaral, A. C. Barros, C. Souza, J. Johns, and J. Gerwing. 1997. Natural resource management in the Brazilian Amazon. *Bioscience* **47**:160-168.

- Uhl, C., and R. Buschbacher. 1985. A disturbing synergism between cattle ranch burning practices and selective tree harvesting in the eastern Amazon. *Biotropica* **17**:265-268
- Uhl, C., and I. C. G. Vieira. 1989. Ecological impacts of selective logging in the Brazilian Amazon: a case study from the Paragominas region of the state of Pará. *Biotropica* **21**:98-106.
- Uhl, C., and J. B. Kauffman. 1990. Deforestation, fire susceptibility, and potential tree responses to fire in the eastern Amazon. *Ecology* **71**:437-449.
- Veríssimo, A., P. Barreto, M. Mattos, R. Tarifa, and C. Uhl. 1992. Logging impacts and prospects for sustainable forest management in an old Amazonian frontier: the case of Paragominas. *Forest Ecology and Management* **55**:169-199.
- Verissimo, A., P. Barreto, R. Tarifa, and C. Uhl. 1995. Extraction of a high-value natural resource in Amazonia: the case of mahogany. *Forest Ecology and Management* **72**:39-60.
- Veríssimo, A., M. A. Cochrane, and C. Souza, Jr. 2002. National forests in the Amazon. *Science* **297**:1478.
- Viña, A., and J. Cavalier. 1999. Deforestation rates (1938-1988) of tropical lowland forests on the Andean foothills of Colombia. *Biotropica* **31**:31-36.
- Walker, G. 1996. Kinder cuts. *New Scientist* **151(2048)**:40-42.
- Whitmore, T. C., 1997. Tropical forest disturbance, disappearance, and species loss. Pages 3-12 in W. F. Laurance and R. O. Bierregaard, editors. *Tropical forest remnants: ecology, management, and conservation of fragmented communities*. University of Chicago Press, Chicago, Illinois.
- Woods, P. 1989. Effects of logging, drought, and fire on tropical forests in Sabah, Malaysia. *Biotropica* **21**:290-298