

**The text that follows is a PREPRINT.
O texto que segue é um PREPRINT.**

Please cite as:

Favor citar como:

Baird, I.G., A.D Ziegler, P.M.

**Fearnside, A. Pineda, G. Sasges, J.
Strube, K.A. Thomas, S. Schmutz,
F. Greimel & D.S. Hayes. nd.**

**Ruin-of-the-rivers? A
global review of run-of-
the-river dams. *Environmental
Management* (in press).**

ISSN: 0364-152X

Copyright: Springer.

The original publication will be available at:
A publicação original estará disponível em:

<https://link.springer.com/journal/267>

Ruin-of-the-Rivers? A Global Review of Run-of-the-River Dams

Ian G. Baird
Department of Geography
University of Wisconsin-Madison
550 N. Park St.
Madison, WI, 53706
USA
ibaird@wisc.edu

Alan D. Ziegler
Faculty of Fisheries Technology and Aquatic Resources
Maejo University,
Chiang Mai
Thailand
ADZ: 0000-0001-5305-2136;
adz@mju.ac.th (preferred: thaihawk@gmail.com)

Philip M. Fearnside
Instituto Nacional de Pesquisas da Amazônia (INPA)
Manaus, Amazonas
Brazil
pmfearn@inpa.gov.br
<mailto:>

Alfonso Pineda
Programa de Pós-graduação em Ecologia de Ambientes Aquáticos Continentais,
Universidade Estadual de Maringá
Brazil
amanecce@gmail.com

Gerard Sasges
Department of Southeast Asian Studies
National University of Singapore
Singapore
GS: 0000-0003-4881-716X;
gerard.sasges@nus.edu.sg

Johann Strube
Territorial Planning Unit
Grand Council Treaty #3
Kenora, ON P9N 3X7
Canada
JohannStrube@cunet.carleton.ca

Kimberley Anh Thomas
Department of Geography and Urban Studies
Temple University
1115 Polett Walk
308 Gladfelter Hall
Philadelphia, PA 19122
USA
kimthomas@temple.edu

Stefan Schmutz
University of Natural Resources and Life Sciences, Vienna, Department of Water, Atmosphere
and Environment, Institute of Hydrobiology and Aquatic Ecosystem Management, Gregor-
Mendel-Straße 33, 1180 Wien
Austria
stefan.schmutz@boku.ac.at

Franz Greimel
University of Natural Resources and Life Sciences, Vienna, Department of Water, Atmosphere
and Environment, Institute of Hydrobiology and Aquatic Ecosystem Management, Gregor-
Mendel-Straße 33, 1180 Wien
Austria
franz.greimel@boku.ac.at

Daniel S. Hayes
University of Natural Resources and Life Sciences, Vienna, Department of Water, Atmosphere
and Environment, Institute of Hydrobiology and Aquatic Ecosystem Management, Gregor-
Mendel-Straße 33, 1180 Wien
Austria
daniel.hayes@boku.ac.at

Abstract

The classification of a hydropower scheme as run-of-the-river (or run-of-river; ROR) evokes an image of a low-impact installation; however, examination of eight case studies worldwide shows that substantial negative societal and ecological impacts are tied to them, albeit in somewhat different ways. We conclude that ROR dams not only potentially displace communities, disrupt livelihoods, and degrade environments in surrounding areas, but they also divert water from areas of need, impact aquatic ecology through habitat destruction and disruption of fish migration, emit non-trivial amounts of greenhouse gases over the lifespan of the project, and disrupt streamflow in downstream river sections. While these negative impacts vary on a case-by-case basis, medium and large ROR dams consistently have multiple and cumulative impacts, even when not having appreciable reservoirs. We contend that many impactful dams do not qualify as low-impact ROR projects, despite being defined as such. Such mislabeling is facilitated in part by the ambiguous definition of the term, which risks the ROR concept being used by proponents of impactful structures to downplay their negative effects and thus mislead the public or gain status, including within the Clean Development Mechanism in relation to mitigating climate change.

Keywords: Run-of-the-river, hydropower, dams, EIA, hydropeaking

1. Introduction

While hydropower has benefited parts of society for more than 150 years (e.g. Von Sperling, 2012; Harrison 2019), their dams have often resulted in myriad negative social and environmental impacts (Baxter 1977; Goldsmith and Hilyard 1984; McCully 2001; Scudder 2005; McManamay et al. 2015). Beyond scholarly critiques, many communities have long raised grave concerns about the negative impacts, lending on-the-ground voices to express mounting worries regarding the integrity of ecologies of local riverscapes and livelihood resources (Goldsmith and Hilyard 1984). Dams constructed on international rivers also remain a tense geopolitical conundrum (Hirsch 2016). A current wave of apprehension focuses on the uncertain role hydropower dams can play in decarbonizing the energy production efforts during a period of climatic uncertainty (cf. Carvajal and Li 2019; Harlan 2020; Sasges and Ziegler 2024).

Since the late 20th Century, the dam-building industry has faced significant resistance to new development from a variety of actors, including local communities, activists, ecologists, and international financiers (McCully 2001; Scudder 2005; Shoemaker and Robichaud 2018). Thus, amidst a period of polarization that pitted proponents of hydropower for economic development against those advocating against it due to environmental costs and social justice concerns (Goodland et al. 1993), the industry began reshaping in an attempt to diminish negative public perception, including advocating for somewhat smaller hydropower facilities (HPFs) and also focusing on improving public relations. This approach also paved the way for the construction of new large dams, albeit with reduced reservoir sizes. This partially but incompletely addressed concerns of large-scale human resettlement, the loss of sensitive or productive lands, and the dramatic disruption of environmental flows in rivers and streams (Cernea 1999; Cernea and McDowell 2000; Scudder 2005; Randell 2022).

Many of these structures were labelled "run-of-river" or "run-of-the-river" (ROR) dams and were often promoted as "environmentally friendly" (Jager and Bevelhimer 2007). Unfortunately, much of this infrastructure has not resulted in the benefits claimed by proponents (Venus et al. 2020).

Contrary to the imagery evoked by the term, ROR dams are not designed to preserve the integrity of all types of environmental flows. Nearly three decades ago, Roberts (1995) used the phrasing “ruin-of-the-river” in reference to 12 proposed mainstream dams of 32 to 46 m height on the Mekong River that would have severe consequences on the aquatic resources and livelihoods of local people, as they are all located at critical locations on the mainstream of the river. These high dam facilities were, and still are, promoted as ROR.

Complementing the rebranding of hydropower dams as ROR is the positioning of hydroelectricity as “green energy” or “climate change friendly energy” (Kahn et al. 2014; Baird and Green 2020). While it may be true that ROR schemes frequently have smaller reservoirs than the early megadams, an important consideration pertains to conjuring the idea that all ROR dams are firstly small, and, secondly, that they are socially and environmentally benign. Recent reviews suggest otherwise (cf. Csiki and Rhoads 2010; Anderson et al. 2015; Kibler and Tullos 2013; Kelly-Richards et al. 2017; Hennig and Harlan 2018; Kuriqi et al. 2021).

This paper challenges prevailing narratives by examining case studies that highlight the significant impacts of ROR dams in Europe, Asia, South America, and North America. The eight case studies presented are based on research conducted by one or more of the authors. These dams vary in size, from 10 MW to over 3,000 MW, and are built on diverse systems, from small mountain streams to large continental rivers. Despite these differences in technology and scale, all are labeled as ROR. The lack of standardized design is a central theme of this paper, as is the ambiguity of the term “run-of-the-river,” which allows it to be used as a rhetorical device to mislead the public, financiers, and government sectors, enabling significant disruption of sensitive riverscapes for profit, often under the guise of development and climate change mitigation. Our goal is not merely to criticize hydropower development but to place our findings within the context of existing research and evaluate whether “run-of-the-river” is a meaningful descriptor for the types of projects that it actually describes.

The paper proceeds as follows. We first briefly present eight case studies of hydropower dams in various parts of the world that developers define as ROR. We then provide a short synthesis of the case studies before briefly reviewing the ROR dam literature related to impacts. We finally provide some concluding remarks.




Figure 1. Map of the locations of case studies included in this paper (source: authors)

2. Case Studies

2.1 FISH MIGRATION DISRUPTION: Pak Mun Dam, Thailand

The Pak Mun Dam in northeastern Thailand exemplifies how medium-sized ROR dams can cause significant environmental and social harm, particularly when situated in locations that obstruct critical seasonal fish migrations (Roberts 1995). In 1967, Thailand's National Energy Office proposed the Pak Mun Dam on the Mun River, a major Mekong tributary in Ubon Ratchathani Province. Initially planned with a large reservoir requiring mass resettlement, it was later redesigned as a smaller ROR dam to reduce displacement. Construction began in 1990 and finished in 1994. The 17 m high dam has a 136 MW capacity, but actual energy output has been lower than expected (Missingham 2003; Foran and Manorum 2009).

Despite its downsizing, the Pak Mun Dam has still caused significant impacts on people forced to resettle due to the creation of the dam's 60 km² reservoir. However, the project's most significant impacts, including in relation to the number of people negatively impacted, have been on seasonal fish migrations between the mainstream Mekong River and the Mun River Basin. Many species migrate up and down the Mekong River during different seasons (Baird et al. 2003; Baran et al. 2005), and the Pak Mun Dam has heavily disrupted many fish migrations that previously migrated upstream from the dam (WCD 2000). Also affected were tributaries of the Mun River that were not originally assessed as affected, and therefore, those impacted have never been compensated (Baird et al. 2020).

The Pak Mun dam's negative impacts on wild capture fisheries have been so severe that since its construction, considerable effort has been devoted to advocating for either opening the dam's gates at certain times of the year or throughout the year to enable fish migration (Foran and Manorum,

2009). Women have been at the forefront of this resistance, adapting older cultural rituals and developing new ones to maintain their opposition to the project (Soukhaphon and Baird 2024). Additionally, the fish ladder added to the dam was poorly designed and has proven ineffective (Roberts 2001).

At one point, the Pak Mun Dam likely stood as the most devastating dam for fisheries ever constructed in the Mekong River Basin, despite being a ROR dam (WCD, 2000). More recently, Ziv et al. (2012) estimated that another highly contentious ROR dam, the Lower Sesan 2 dam in northeastern Cambodia, would become the most damaging dam to fish ecology and fisheries in the Mekong Basin. This project is situated along a crucial migratory fish passage. These two projects highlight that ROR dams can be devastating when it comes to obstructing vital fish migrations.

2.2 BIODIVERSITY: Danube Salmon

Three proposed small-scale (<10 MW) ROR hydropower projects planned for the Mur River in Central Europe are a grave threat for Danube salmon (*Hucho hucho*), the world's largest salmonid species, which is endemic of the Danube River basin (Holčík et al. 1988) and is symbolic of the region's ecological heritage. In recent decades the salmon have faced extensive habitat loss and hydropower-related impacts (Schmutz et al. 2002). Once widespread, this flagship species is now one of the most threatened species in the catchment (Freyhof and Kottelat 2008) and is protected under European conservation law (Annex II and V of the EU Habitats Directive - 92/43/EEC). Recent assessments in Bavaria, Germany, and Austria (Schmutz et al. 2023) revealed that merely 0.7% of its original 7,500 river kilometers habitat range still hosts very good populations, with most rivers showing only moderate to poor population status, reflecting fragmented habitats and dwindling reproduction rates (Schmutz et al. 2023). This situation underscores the urgency of safeguarding the Danube salmon.

In Austria, the last bastions of Danube salmon persist in three (sub-) catchments: the Mur, Gail, and Pielach Rivers. These fish, known for their migratory behavior and preference for gravelly substrates, are critically dependent on free-flowing rivers. Notably, the few remaining (very) good Danube salmon populations in Austria are mainly located in rivers with longer free-flowing sections (>50 km), defined as river sections without dams, impoundments, and flow alterations (water abstraction, hydropeaking), such as the Mur and Gail Rivers (Figure 2). In addition, Danube salmon stocks in the mainstem Danube and lower Mur River suffer from increased summer temperatures. These pressures have led to severe declines in Danube salmon stocks in Austrian rivers (Schmutz et al., 2023), which have been further compounded by losses in prey species such as European grayling (*Thymallus thymallus*) (Hayes et al. 2021), nase (*Chondrostoma nasus*), and barbel (*Barbus barbus*) (Hayes et al. 2022a).




Figure 2. Former and present distribution of Danube salmon in Austria, remaining rivers with free-flowing river sections (>50 km), and the planned hydropower plants at the Mur River (data source: Schmutz et al., 2023).

Despite generating a mere 0.3% (120 GWh) of Austria's annual hydropower output, the three proposed ROR projects pose a significant risk to the last upper Danube River stretches harboring robust Danube salmon populations. Alarming, some of these projects are situated within Natura 2000 protected areas. Despite environmental campaigns against them, politicians have largely supported their development plans are largely supported by politicians, and Europe's emergency council regulation (EC 2022/2577) has also supported them in their push for more renewable energy.

This case study underscores the critical need to scrutinize ROR hydropower expansion, considering the ongoing biodiversity crisis and the imperative to safeguard this irreplaceable species for future generations, particularly given the limited energy benefits provided by these relatively small-scale projects.

2.3 HYDROPEAKING: Austria-wide survey

This case study explores the large-scale potential of ROR hydropower to cause sub-daily flow fluctuations (Almeida et al. 2020), a phenomenon commonly associated with storage hydropower (i.e., hydropeaking), and which cause severe impacts on river ecosystems (Hayes et al. 2022b; Schmutz et al. 2015). To this aim, sites situated within Austrian mountain rivers (Hayes et al. 2021) were categorized into three groups based on their peaking frequency (Figure 3a): low (<0.2 daily peaks), medium (0.2–2.1 daily peaks), and high (>2.26 daily peaks). River sections experiencing medium to high levels of flow fluctuation frequency were predominantly found near peak-operating hydropower plants (Figure 3b). Intriguingly, some negatively affected sites were discovered close to ROR hydropower plants, even when no peak-operating storage dam was present in the upstream catchment, lending evidence that ROR facilities can contribute to sub-daily

flow fluctuations. However, this is not always the case, as evidenced by sites classified under the 'low' category (Figure 3c).




Figure 3. Hydro-ecological effects of artificial flow fluctuations at Austrian fish sampling sites ($n=69$; Hayes et al., 2021). This figure categorizes the sites into 'low', 'medium', and 'high' groups based on (a) the peaking frequency terciles. Panels (b) and (c) illustrate the proximity of each site to the nearest hydropeaking (HP) or ROR power plant, respectively. Panels (d) and (e) present ecologically relevant flow fluctuation metrics (amplitude and down-ramping rate), each normalized against the long-term mean annual maximum flow event (Greimel et al., 2016; Greimel, 2022). Panel (f) depicts the response of the 'Fish Index Austria', a fish-based measure of ecological integrity (1 being best, 5 worst). Group differences were determined with the Kruskal-Wallis (KW) test, followed by pairwise comparisons with the Mann-Whitney U test.

Sites characterized by having a low peaking frequency experienced events with the highest amplitudes compared to those in the medium and high frequency groups. These significant flow fluctuations predominantly constitute natural floods. Conversely, the peak amplitudes observed in the medium and high frequency groups are significantly lower (Figure 3d), which is likely due to the maximum turbine flow of the hydropower plants and a corresponding limitation of the artificial flow fluctuation amplitude. Sites experiencing medium to high event frequencies are distinguished by rapid down-ramping rates (Figure 3e), which can have significant ecological impacts, including fish stranding (Hayes et al. 2022b; Moreira et al. 2019). These hydrological pressures result in a

decline in ecological integrity, as evidenced by the scores from the multi-metric ‘Fish Index Austria’, indicating step-wise degradation (Figure 3f).

This study on Austrian rivers reveals that ROR hydropower installations often produce a high frequency of sub-daily flow fluctuation events, like traditional hydropower plants with large storage. Others have indicated that ROR installations can cause such flow variations, for example, due to delayed reactions of the power plant flow control, low flow stoppages, system start-ups, powerhouse breakdowns, intake malfunctions, cycling operations, and forebay oscillations (Greimel et al. 2016; Hunter 1992). While these artificial fluctuations have a smaller amplitude than natural floods, they occur frequently and exhibit a fast down-ramping intensity, thus causing significant negative impacts on river ecology.

2.4 TRANSBASIN WATER TRANSFER: Theun-Hinboun Dam, Laos

The Theun-Hinboun Dam in Central Laos illustrates how some projects are classified as ROR despite having nothing to do with maintaining natural flow regimes. The Theun-Hinboun Hydropower Project, which is located in Bolikhamxay and Khammouane Provinces in the Lao People’s Democratic Republic (Lao PDR), originally had a 220 MW installed capacity when it started operations in 1998. In 2012 the project was expanded to 500 MW capacity. The dam was developed in partnership between the Lao government, Thailand’s MDX Lao Public Company, and Nordic Hydropower AB, with considerable loans to the Lao PDR government from the Asian Development Bank.

The dam was built based on a 30-year “build–operate–transfer” (BOT) agreement with a 10-year optional extension period, after which the dam would become fully owned by the Lao government (Shoemaker 1998; Whittington 2018). The original project included a large 48 m high dam across the Theun River, a 49 km² reservoir, and infrastructure to facilitate the diversion of water from the dam’s reservoir through a power generation plant and then into the Hai and Hinboun Rivers, tributaries to the Mekong River. The expansion was agreed to in 2007, and involves a 65 m high dam, and a 105 km² reservoir on the Gnouang River.

As various studies have indicated, the dam has led to a significant increase in water flow down the Hai and Hinboun Rivers to the Mekong, sometimes more than double the previous levels (Barney 2007; Blake et al. 2005; Shoemaker 1998). This diversion has resulted in well-documented downstream social and environmental impacts, including significant losses and long-term consequences for downstream communities, such as riverbank erosion, the destruction of aquatic habitat, and various impacts on human livelihoods, such as the loss of riverbank gardening and dramatic declines in fish catches (Barney 2007; Blake et al. 2005; International Rivers Network 1999; Shoemaker 1998; Warren 1999; Whittington 2018). In contrast, the reduction of flows in the Theun River has resulted in substantial social and environmental impacts downstream on the Theun (Shoemaker 1998; Warren 1999).

This example highlights a discrepancy in the application of the term ROR to describe the flow situation on the river where the facility was built. A crucial aspect is the diversion of water from one river (the Theun) to others (the Hai and Hinboun), is the categorizing of the diversion project as a “transbasin run-of-the-river project” (Shoemaker 1998; Warren 1999). The question arises as to whether a dam can be accurately classified as “run-of-the-river” when it redirects water from

one river into another. The Theun Hinboun Power Company argues that it can, because the dam does not have significant active storage. This indicates how the term "run-of-the-river," at least to some actors, does not imply that the river is allowed to flow naturally. Instead, it is a technical term that relates to a dam's active storage. In particular, this reveals how the "politics of classification" are employed to divert attention from a project's serious negative impacts.

2.5 DISPLACEMENT: Belo Monte Dam Complex, Brazil

Brazil's Belo Monte complex illustrates the falseness of ROR dams being either small, or "low impact." Belo Monte is classified as ROR because the water level in the reservoir only fluctuates by one meter, making the complex depend on the flow of the river. The complex began blocking the Xingu River, a north-flowing tributary to the Amazon River, in 2015 (Figure 4). It was originally planned that up to five large storage dams upstream would have regulated flow, which would have flooded vast areas of Indigenous land (Fearnside 2006). Even though Brazil's official statements since 2008 have claimed that the upstream dams will not be built, turbines with installed capacity totaling 11,233 MW have already been installed at Belo Monte (an amount around twice the maximum that could be justified by the river's natural flow), suggesting that plans for the upstream dams continue unannounced (Fearnside 2017a). Since 2013, when the dam was under construction, Belo Monte has been used by Brazil's authorities to argue for future projects that prioritize water storage (Fearnside 2017b).




Figure 4. Map of the Belo Monte complex and surrounding area (Fearnside, 2017a).

The Belo Monte complex consists of two dams: the Pimental Dam functions as an intra-basin water diversion scheme. It diverts 80% of the river's flow through a canal and a series of dammed stream basins to a second dam (Belo Monte). The operational mode leaves a 130 km long section of the river with only 20-30% of its natural flow. Indigenous communities on the banks of this section of the Xingu River, known as the “*Volta Grande*” (or “Big Bend”), lost the fish and turtles that previously sustained them, as did a third group located on the Bacajá River, a tributary (Pezzuti et al. 2024).

The 516 km² reservoir created by the dam displaced 40,000 people, according to the Movement for Dam Affected People (Sullivan 2016), including both traditional riverside residents (*ribeirinhos*) and one-fourth of the city of Altamira. The displaced people were moved to “collective urban resettlements” on the outskirts of Altamira, where they had no income or means of producing food. *Ribeirinhos* and the Indigenous Peoples who remained along the *Volta Grande* lost almost all fish and other resources that sustained them (Magalhães and da Cunha 2017).

Besides displacing people, the Pimental Dam also blocked the migration of fish, despite having a fish passage, as most Amazonian fish fail to pass these barriers. Both the area flooded by the reservoir and the *Volta Grande* had rich endemic fauna that has now been severely damaged (Keppeler et al. 2022). The most famous victim is the zebra pleco (*Hypancistrus zebra*), a valuable aquarium fish adapted to rocky habitats in rapids. This species is expected to go extinct in the wild (Gonçalves 2011). The rapid variations in water level have taken a heavy toll on commercial fish species during the spawning season because they enter flooded forest areas that suddenly become either dry land or isolated pools of intolerably hot water (Ribas et al. 2023).

The Belo Monte complex also accelerated deforestation (Jiang et al. 2018) by attracting people to the area and increasing land values and yielding profits from land speculation (Barreto et al. 2011). Furthermore, a study measuring greenhouse gas (GHG) emissions at Belo Monte concluded that “...total GHG emissions are substantial even from this leading-edge ROR power plant. This argues in favor of avoiding hydropower expansion in Amazonia regardless of the reservoir type” (Bertassoli Jr. et al. 2021). The Science Panel for the Amazon concluded that no more dams of any type with installed capacity ≥ 10 MW should be built in Amazonia (Fearnside et al., 2021), as even dams smaller than 10 MW can cause significant impacts depending on various factors.

It may be a surprise to many that ROR dams are sometimes quite large and can result in large-scale human resettlement as well as many of same sorts of impacts associated with large storage dams.

2.6 LIVELIHOOD DISRUPTION: Rainy River International Dam, USA-Canada border

The 24.4 MW International Dam spans the Rainy River between Minnesota (USA) and Ontario (Canada). Because of the area's flat topography, this comparably small dam (9.1 m height) maintains the water level in the 932 km² Rainy Lake (*Gojiji-zaaga'igan*), a former wetland turned reservoir. European-descended residents and seasonal tourists typically only pay attention to the dam in high water years when lake water could have been released sooner to avoid flooding (Kraker 2022; IRLWWB 2023). However, the Indigenous Anishinaabeg in the area perceive the dam differently.

Prior to the construction of the dam between 1905 and 1909, the marshes above the dam sustained abundant Manoomin (wild rice, *Zizania palustris*) crops that sustained Indigenous Peoples for millennia (Birk and Richner 2004; Reid and Rajnovich 1991). Pushed by the newsprint industry and enabling settler-colonial governments, construction of the dam and others in the area virtually destroyed the Anishinaabe wild rice economy. *Manominikenshii*, the Anishinaabe wild rice culture, evolved under specific hydrological conditions regarding seasonally variable water levels, temperature, pH, turbidity, and nutrient loads affected by natural conditions, localized management (e.g., impoundment of bays and streams, seeding), and Ceremony (Strube 2021; Kinew 1995; Waisberg 1984). The dam, despite its characterization as a small ROR facility and temporary impoundment, has disrupted this delicate ecological system, upsetting the Anishinaabe's relationship with Manoomin with grave consequences for the Anishinaabeg's livelihood and sovereignty (Strube and Thomas 2021). Anishinaabe communities on the lake's southern shore migrated south due to the dam's effects on their economy (Child 2011). Without their main food staple, the Anishinaabeg on the Northern shore lost much of their autonomy and became increasingly dependent on the settler-colonial state, which in turn expanded its reach into this borderland through industrialization dependent on hydropower and water-level regulation (Strube 2021; Strube and Thomas 2021).

That a dam with such far-reaching impacts can be perceived by settlers as innocuous is the result of several discursive moves. Most critically, the hydrological studies and deliberations informing water regulation on the Rainy River have long erased Indigenous Peoples as political agents and rightsholders. Consequently, dam operations have long ignored wild rice and instead privilege the interests of waterfront residents, tourism, and local industry by providing constant lake levels.

Today, the facility is depicted by the operator as a “run-of-the-river plant” that is subject to restrictions on peaking operations, including adherence to Rainy Lake rule curves, minimum flow requirements, and control orders set by the International Joint Commission. However, while the dam does not retain and release water on demand like storage dams typically do, it does hold back water for “a few days,” as one of the operating engineers admitted. Contradicting their public relations department's framing, an engineer called into question the characterization of the dam as ROR, instead offering “intermediate semi-run-of-river” as an alternative, yet similarly misleading classification (Strube 2021). This case study shows that reimagining older dams as ROR projects focuses public attention on environmental flows that align with contemporary management protocols, meanwhile detracting attention from significant negative impacts on local communities.

2.7 GREENWASHING: Đăk Psi 3 and 4 cascade, Central Vietnam

Despite its relatively short length of 81 km, the Đăk Psi River in Central Vietnam's Kon Tum province is the site of ten hydropower facilities on its main stem and tributaries. The Đăk Psi 3 (15 MW) and 4 (30 MW) cascade (hereafter DP3-4) was the first constructed, from 2007 to 2012. The project was subsidized through the Clean Development Mechanism (CDM) of the United Nations, which grants certified emission reduction credits to corporations that invest in renewable energy projects (Martins et al. 2013; Erlewein 2014). As part of CDM certification, the project was designated as ROR, which from the perspective of Vietnamese officials, implies minimal environmental impacts.

CDM-related and environmental impact assessment (EIA) reports consistently refer to the project as ROR; and diagrams depict the construction of “weirs” without reservoirs, allowing for continuous flow. However, as built, the weirs create reservoirs and obstruct the flow of water, materials, and organisms (Figure 5). The claim that the project is ROR rested on the use of 2-3 km-long diversion tunnels leading from the weirs to powerhouses from where water was discharged back into the river channel.




Figure 5. (A) Redrawing of the schematic of the Đak Psi 3-4 project submitted for EIA and CDM approval, circa 2011 (Dak Psi, 2023; the original orientation and features/elements are maintained, except those highlighted in red). (B) Đak Psi 3-4 cascade circa 2021. Structures labelled “weirs” function as dams, creating 15-25 ha reservoirs and limiting downstream flows along a length of 2-3 km.

ROR labeling was crucial both to securing CDM financing and a positive EIA. The CDM Project Design Document states the following (UNFCCC 2006): “Đak Psi 3, 4 Hydropower Project is a run-of-river type. ... Therefore, the negative impacts are fairly limited in scope.” In turn, the EIA and consultative process carried out prior to the dam’s construction proceeded from the assumption that a small ROR facility would have limited environmental impact. It did not highlight some of the most important foreseeable impacts of the facility, including the blocking of environmental flows, ecological fragmentation, and the dewatering of downriver reservoirs. Impact assessments focused on short-term effects during construction rather than the long-term impacts on local ecologies.

Contrary to the scenario sketched out by project backers, the years since the construction of the dams along the Đak Psi have seen the transformation of the river’s ecologies. After completion, the stream section below the dam became dewatered. Villagers describe how, before the dam, the riverbanks were rocky and supported lush green trees. Today, due to increased sedimentation downstream, sand has filled up the riverbed, raising the water and exposing farmland along the riverbank to floods during the wet season. In addition, during the dry season, water levels in the river are insufficient for their daily human needs (Du Toan 2022).

In this case, the transformation of local lives, livelihoods, and ecologies was justified in the name of the decarbonization of energy production. CDM financing is intended to support green, carbon-offsetting projects that would not otherwise be built due to a lack of capital. However, the construction of Đak Psi 4 began in 2007, one year before the project's operators applied for CDM financing. Thus, the carbon credits associated with the CDM financing effectively subsidized further fossil fuel use, resulting in a net gain in carbon emissions and implicating DP3-4 in a global system of "greenwashing" of hydropower, which has been termed the "theater of decarbonization" (Sasges and Ziegler 2023; 2024). The DP3-4 cascade case study demonstrates how manipulation of terms supports greenwashing, silencing critiques of hydropower project impacts. This is part of a global rhetoric exploiting the idea that all hydropower, regardless of its specifics, helps reduce climate change.

2.8 CUMULATIVE IMPACTS: Jirau and San Antonio Dams, Brazil

Brazil's Santo Antônio and Jirau hydroelectric plants on the Madeira River, a key Amazon tributary, were completed in November 2011 and September 2012, respectively. These are the first dams in the Madeira River complex. Both projects contribute to the regional power grid, with installed capacities of 3,580 MW (Santo Antônio) and 3,750 MW (Jirau). Being very large dams at 49.5 m (Santo Antônio) and 62 m (Jirau), it is not surprising that grave concerns were raised in their Environmental Impact Report (RIMA) and EIA. The environmental agency also faced intense political pressure to approve licensing (Fearnside 2014a). These dams are classified as ROR because there is very little fluctuation of the water level behind them, and because power is being generated by the river's natural flow rather than by drawing down stored water (Fearnside, 2014a). During the licensing process, documents such as the strategic environmental evaluation and the economic viability study prominently featured a photograph of a ROR dam with no reservoir, located on Europe's Danube River, falsely implying that the Madeira River dams would be similarly reservoir free (Fearnside 2014b).

The dams in the Madeira River have disrupted the distribution of fish throughout the basin, which, in turn, has impacted the livelihoods and food security of over 50,000 people across Peru, Bolivia, and Brazil (Doria et al. 2018). The Santo Antônio and Jirau Dams have negatively affected the migration of the large catfish, *Brachyplatystoma rousseauxii*, known as "dourado," which, prior to the dams, was responsible for generating over 14,000 tons of fish biomass just in the Brazilian portion of the river. *Dourado* migrated from the mouth of the Amazon River and ascended the Madeira River to reproduce in the Bolivian and Peruvian Andes, and larvae and juveniles descended to the Amazon River's mouth. The Santo Antônio and Jirau Dams now obstruct the migration of both adult and juvenile fish (Hauser et al. 2019).

Dam construction raised the Madeira River's average water level by 10 m, reshaping the landscape. The flooding associated with the complex created backwater zones at tributary junctions, causing chemical changes and thermal stratification. The flooded area following the installation of the Madeira complex exceeded the planned area by 341 km² (64.5%), submerging an additional 160 km² of natural forest (Cochrane et al. 2017). These losses have had profound social consequences, displacing thousands of people from their homes, destroying their livelihoods, and significantly reducing their access to food (Fearnside 2014a).

The decomposition of inundated plant biomass, coupled with the formation of backwater regions, introduces an additional concern related to the Jirau and Santo Antônio Dams: the release of greenhouse gases. Dams in the Amazon region emit methane (CH₄) and carbon dioxide (CO₂) from both the vegetation and the soils that are flooded, and the emissions from the flooded vegetation elevate CH₄ emissions by 33% and CO₂ emissions by 28%, compared to the soil alone (de Faria et al. 2015). Collectively, the Jirau and Santo Antônio Dams will emit an estimated average of 76 Tg of carbon over 100 years, equivalent to nearly 800,000 tons of carbon annually.

These dams received carbon credits through the CDM (see Vietnam case study above) despite having been built for reasons unrelated to carbon reduction, thereby allowing the countries that purchased these credits to emit without really offsetting carbon emission, thus contributing even more to global warming (Fearnside 2015). Carbon projects often assume hydropower dams have no emissions; however, methane generated in stratified tributaries entering the Santo Antônio reservoir has been found to be released in substantial quantities downstream of the dam (Fearnside, 2015). These findings underscore the fact that ROR projects can have significant environmental and social impacts, including cumulative impacts, both locally and regionally, as well as significantly contributing to greenhouse gas emissions.

4. Discussion

4.1 Synthesis of case study impacts

The case studies in Section 3 reveal how the term "run-of-the-river" (ROR) has been inconsistently applied to hydropower facilities of varying designs and scales. In addition, the labeling of these dams as ROR by developers often fails to reflect their significant environmental and social impacts. For instance, the construction of Brazil's Belo Monte Complex resulted in the displacement of approximately 40,000 people, highlighting severe social consequences akin to those criticized in mega dam projects. Similarly, the Santo Antônio and Jirau Dams on the Madeira River, with heights of 50-60 m, created reservoirs nearly two-thirds larger than anticipated, flooding 16,000 hectares of natural forest beyond the predicted area.

Damming the Rainy River has led to the loss of wetland habitats long crucial to the Indigenous Anishinaabe peoples, echoing a broader trend of livelihood disruptions caused by large ROR dams. Such impacts extend to other groups affected by dams such as Pak Mun in Thailand and Theun-Hinboun in Laos, where hydrological changes and other ecological impacts have been significant.

Ecologically, ROR dams can severely disrupt fish migrations, fragment fish populations, and damage critical habitats, as seen with the Pak Mun Dam in Thailand and the series of ROR dams along Austria's Mur River. A particularly concerning issue is turbine mortality for downstream-migrating fish, even with mitigation measures like bar racks and migration corridors. This mortality, especially for juvenile and small-bodied fish, varies by species and turbine type, but the cumulative effects from multiple passages can drastically reduce fish populations (Knott et al. 2023a, b; Pracheil et al. 2016; Radinger et al. 2022).

Although ROR dams have often been associated with maintaining environmental flows or replicating natural downstream patterns (Dyson et al. 2003; Hayes et al. 2018), the case studies presented here demonstrate that water diversions and hydropeaking can significantly affect downstream flow regimes. For example, the Theun-Hinboun Dam in Laos diverts substantial water flow between river catchments, and Brazil's Belo Monte Dam dramatically reduces flows over a

130 km stretch through diversion. Similarly, the Dak Psi 3-4 cascade in Vietnam diverts water from the dam to the power plant 2-3 km downstream, leaving the intermediate river reach dry for much of the year. In Austria, ROR dams have also caused significant downstream impacts through daily hydropeaking, despite lacking the active storage typically associated with such effects.

Some projects have used the ROR label for greenwashing purposes, including securing environmental approvals and funding, despite causing considerable harm. For instance, the Jirau and Santo Antônio dams in Brazil and the Đak Psi 3/4 dams in Vietnam were branded as ROR to gain credibility under the CDM, even as they resulted in significant environmental and social impacts that were often downplayed or ignored in assessments (Baird and Green, 2020). Ironically, these dams also emit substantial greenhouse gases, contradicting the CDM's intentions (Barros et al. 2011).

4.2. Other Evidence of Adverse ROR Impacts

Complementing the impacts reviewed for the eight case studies in Section 3 are numerous investigations conducted worldwide. We found 59 empirical studies on ROR impacts using a systematic search on related keywords. These studies are from 20 countries, including Austria, Brazil, Canada, China, France, India, Japan, Norway, and the USA. The ROR facilities range from small weirs to large dams on international rivers such as the Danube, Rhône, and Columbia. This diversity reflects the lack of a common definition or design for ROR dams globally (Csiki and Rhoads 2010; McManamay et al. 2016; Kuriqi et al. 2021).

Many studies focus on specific impacts, such as the adverse effects on benthic organism communities, including macroinvertebrates (Dessaix et al. 1995; Fanny et al. 2013; Wang et al. 2016; Anderson et al. 2017; Bilotta et al. 2016; Silverthorn et al. 2018; Mihara et al. 2024). Other organisms affected include diatoms (Wu et al. 2010; Wang et al. 2022a, b), algae (Shuka et al. 2013; Wu et al. 2009), plankton (Li et al. 2018; Rose et al. 2019; Zanon et al. 2024), amphibians (Dare et al. 2020), and insects (Malmqvist and Englund 1996). Negative impacts on fish community dynamics are also well-documented (Simonović et al. 2021; Baumgartner et al. 2020; Ticiani et al. 2023).

For instance, on the Tay River in Scotland, Robson (2013) observe that ROR dams reduced water flow in depleted reaches, restricted upstream access, and caused combined barrier and abstraction effects. Linares et al. (2018) report that ROR impacts primarily affected dam reservoirs and adjacent downstream stretches, facilitating invasive species dominance. Bejarano et al. (2019) note that ROR-diversion power plants impact river reaches downstream, particularly in nival and stable river types. Magilligan et al. (2021) observe that while ROR dams do not significantly disrupt sediment connectivity, they affect ecological connectivity.

Research on the St. Maurice River in Canada shows that ROR dams can disrupt ecosystem processes such as carbon cycling and mercury dynamics, potentially leading to elevated mercury levels in aquatic organisms (Ponton et al. 2021; Leclerc et al. 2023). Tashiro et al. (2015) report that ROR dams can reduce daily gross production and community respiration below dams under low flow conditions. Almeida et al. (2019) documents increased biochemical oxygen demand and CO₂ partial pressure due to the influx and mineralization of organic matter. Sow et al. (2016) highlight the role of shallow depths and submerged macrophytes in nutrient and sediment retention.

Several studies highlight hydro-geomorphological impacts. Reduced stream flow can lead to sediment accumulation and erosion downstream due to sediment shortages and increased flood velocities (Summer et al., 1994). For a large ROR facility on the Rhône River, significant geomorphic changes due to a century-long reduction in sediment supply have been observed (Dépret et al. 2019). Other research indicates that low-head ROR dams create conditions of limited sediment supply downstream, affecting local and downstream ecological habitats (Casserly et al. 2020).

In Italy, the River Po experienced significant degradation following ROR dam construction due to altered flow regimes and sediment transport disconnection (Bizzi et al. 2015). In Spain, the Upper Garonne River saw channel narrowing after ROR dam construction, leading to new management practices involving downstream flushing actions (Bulteau et al. 2022). Wildman and MacBroom (2005) report that low-head ROR dams cause sediment accumulation and stream widening, leading to dam removal. Pearson et al. (2016) note that ROR dams may induce brief periods of sediment methane flux to the atmosphere, with potential impacts from dam removal.

From a social perspective, Rousseau (2020) report on greenwashing tactics in Yunnan, China, where a dam recognized as ROR caused significant impacts to local villagers. Ullah et al. (2023) note that a large ROR plant on the Indus River in Pakistan affected river flow regimes, groundwater levels, and land use, exacerbated by inadequate compensatory measures.

In contrast, some studies report minor impacts. Csiki and Rhoads (2014) found that small ROR dams in Illinois do not substantially alter channel morphology or act as major sediment traps. Hocking et al. (2021) report an increase in rainbow trout biomass due to controlled flow diversions, maintaining natural flow regimes. Copeman (1997) observe no adverse effects on sediment and benthic invertebrates but cautioned about rivers with more variable flows. Neupane et al. (2023) report a favorable increase in ecosystem service values associated with ROR development in Nepal, despite significant land use changes.

Finally, in support of the studies from the literature, several syntheses conclude that ROR hydropower plants have moderate to significant impacts on river ecosystems. Anderson et al. (2015) conclude that ROR schemes can reduce habitat complexity, alter riparian vegetation, and disrupt longitudinal connectivity. Csiki and Rhoads (2010) discuss how ROR dams impact river geomorphology, potentially causing downstream scour. Gibeau et al. (2017) identify three pathways through which ROR hydropower affects salmonids. Kuriqi et al. (2021) emphasize that small ROR plants significantly alter natural flow regimes and harm fluvial ecosystems.

4.3. Prospects for Monitoring and Adaptive Management

The above sections describe socio-ecological impacts of ROR facilities, impacts that are often much more seriously than typically recognized. However, a certain degree of mitigation is possible. Best-practice mitigation measures often related to the operation of these projects. For example, water can be released to more closely mimic natural flows (an environmental flows regime). This can include periodic channel-forming discharges (Hayes et al. 2018), water releases for the purposes of fish protection (Haug et al, 2022). In addition, fish passes to support

migration (Silva et al., 2018) and restrictions on flow ramping (hydropeaking) (Moreira et al., 2019) can also help partially mitigate the impacts.

However, a strict monitoring scheme is needed at existing facilities to ensure that projects are managed to reduce environmental impacts. This needs to include regular assessments of water quality (including water temperature) and aquatic biota, but also continuous monitoring of river discharge. Involving local people and making data publicly available is also important to help ensure that impacts are minimized. The iterative framework of adaptive management can help to formulate new management actions by integrating new information and paying close attention to both ecological and social factors (Gunderson et al., 2016; Sendzimir & Schmutz, 2018). Ultimately, however, we contend that many of the serious negative impacts of ROR dams cannot be easily or fully mitigated.

5. Conclusion

The review of eight case studies, along with numerous studies from the literature consulted, reveal that while ROR dams may appear less harmful than large storage dams, they frequently result in significant negative impacts, including community displacement, ecological degradation, and disruption of local livelihoods. This reality challenges the prevailing perception that ROR dams are benign sources of renewable energy.

We identify substantial inconsistencies in how the term "run-of-the-river" is applied across different geographical, ecological, and socio-political contexts. ROR is broadly used to describe various hydraulic structures built across river channels for hydropower generation, but its definition lacks clarity. For some, ROR refers to dams that form reservoirs without active storage or have insufficient capacity for seasonal water management. Others use the term to denote dams that maintain minimal discharge to preserve environmental flows, or they associate ROR with small hydropower facilities, regardless of their specific design. The absence of a standardized definition renders the term ROR susceptible to rhetorical manipulation rather than being a precise engineering concept. This ambiguity allows hydropower proponents to downplay the potential negative impacts of ROR dams. As Csiki and Rhoads (2010) argue, ROR is not a scientific term but a commonly accepted phrase in river management, likely derived from older "reservoirless" hydroelectric facilities.

Historically, hydropower advocates have used ROR to describe dams with minimal dead storage needed for consistent water release to generate electricity. Despite claims that ROR dams allow rivers to flow freely, their designs frequently prioritize optimizing water release for power generation over maintaining natural river dynamics (Anderson et al. 2015). Further, definitions that describe ROR as systems where discharges approximate the sum of inflows—relying on natural flows and minimal reservoir fluctuations—often neglect the impact of structures that obstruct two-way flows, significantly disrupting migrating fish and other aquatic organisms. While ROR dams often feature smaller reservoirs than traditional storage dams, which may result in less severe hydrological alterations and fewer resettlement issues, they can still cause substantial disruption to social and ecological systems. This aligns with early warnings from Roberts (1995) regarding the detrimental impacts of ROR dams proposed to be built on the Mekong River.

In conclusion, the numerous reports of negative impacts of ROR dams challenge the narrative that they are inherently clean and sustainable, underscoring the need for a more nuanced understanding of ROR facilities. To address this issue, it may be necessary to reserve the term ROR for designs that genuinely preserve environmental flows and develop new classifications that more accurately reflect the disruptive features of other hydropower facilities. Such measures are essential not only to prevent the misleading use of the term ROR but also to foster hydropower development as a genuinely renewable energy source that benefits society as a whole.

References

- Almeida RM, Hamilton SK et al. (2019) Limnological effects of a large Amazonian run-of-river dam on the main river and drowned tributary valleys. *Scientific Reports* 9(1):16846.
- Almeida RM, Hamilton SK et al. (2020) Hydropeaking operations of two run-of-river mega-dams alter downstream hydrology of the largest Amazon tributary. *Front in Environ Sci* 8. <https://doi.org/10.3389/fenvs.2020.00120>.
- Anderson D, Moggridge H et al. (2017) Quantifying the impact of water abstraction for low head ‘run of the river’ hydropower on localized river channel hydraulics and benthic macroinvertebrates. *River Res and Applic* 33(2):202-213.
- Anderson D, Moggridge H et al. (2015) The impacts of ‘run-of-river’ hydropower on the physical and ecological condition of rivers. *Water and Enviro Jour* 29(2):268-276. <https://doi.org/10.1111/wej.12101>.
- Baird IG, Green WN (2020) The Clean Development Mechanism and large dam development: contradictions associated with climate financing in Cambodia. *Clim Change* 161(2):365-383.
- Baird IG, Manorum K Phenow et al. (2020) Opening the gates of the Pak Mun dam: Fish migrations, domestic water supply, irrigation projects and politics. *Water Altern* 13(1):141-159.
- Baird IG, Flaherty MS and Phylavanh B (2003) Rhythms of the river: Lunar phases and migrations of small carps (Cyprinidae) in the Mekong River. *Nat Hist Bull Siam Soc* 51(1):5-36.
- Baran E, Baird IG, Cans G (2005) Fisheries bioecology in the Khone Falls area (Mekong River, Southern Laos). In Baran E, Baird IG, Cans G (Eds), *Bioecology of Khone Falls Fisheries (Mekong River, Southern Laos)*. WorldFish Center, Penang, Malaysia.
- Barney K (2007) Power, progress and impoverishment: plantations, hydropower, ecological change and community transformation in Hinboun District, Lao PDR. *Political Ecology Series, (Re)Making Governance Series, YCAR Papers #1*, York University, Toronto, ON.
- Barreto P, Brandão Jr. A et al. (2011) Risco de desmatamento associado à hidrelétrica de Belo Monte. Instituto do Homem e Meio Ambiente da Amazônia (IMAZON), Belém, PA, Brazil. 98 pp. <https://bit.ly/47O3tYL>.
- Barros N, Cole JJ et al. (2011) Carbon emission from hydroelectric reservoirs linked to reservoir age and latitude. *Nature Geosci* 4(9):593–596. <https://doi.org/10.1038/ngeo1211>.
- Baumgartner MT, Piana PA et al. (2020) Storage or run-of-river reservoirs: exploring the ecological effects of dam operation on stability and species interactions of fish assemblages. *Environ Manag* 65(2):220-231.
- Baxter RM (1977) Environmental effects of dams and impoundments. *Annual Review of Ecology and Systematics* 8(1):255-283. <https://doi.org/10.1146/annurev.es.08.110177.001351>.

- Bejarano MD, Sordo-Ward A et al. (2019) Tradeoff between economic and environmental costs and benefits of hydropower production at run-of-river-diversion schemes under different environmental flows scenarios. *Jour of Hydro*. doi:10.1016/j.jhydrol.2019.03.048.
- Bertassoli Jr DJ, Sawakuchi HO et al (2021) How green can Amazon hydropower be? Net carbon emission from the largest hydropower plant in Amazonia. *Sci Adv* 7, art. eabe 1470. <https://doi.org/10.1126/sciadv.abe1470>.
- Bilotta GS, Burnside NG, et al. (2016) The effects of run-of-river hydroelectric power schemes on fish community composition in temperate streams and rivers. *PLoS One* 1–15. doi:10.1371/journal.pone.0154271.
- Birk DA, Richner J (2004) *From Things Left Behind: A Study of Selected Fur Trade Sites and Artifacts, Voyageurs National Park and Environs, 2001-2002*. Voyageurs National Park, International Falls, MN.
- Blake D, Carson B, Tubtim N (2005) *Review of the Environmental Management Division*. Theun-Hinboun Power Company, Vientiane.
- Bulteau T, Batalla RJ et al. (2022) Geomorphic effects of a run-of-the-river dam in a multi-driver context: The case of the Upper Garonne (Central Pyrenees). *Geomorphology* 408:108243.
- Carvajal PE, Li FG (2019) Challenges for hydropower-based nationally determined contributions: a case study for Ecuador. *Clim Pol* 19(8):974-987. <https://doi.org/10.1080/14693062.2019.1617667>.
- Cassery CM, Turner JN et al. (2020) Impact of low-head dams on bedload transport rates in coarse-bedded streams. *Sci of the Total Environ* 716:136908.
- Cernea MM (Ed) (1999) *The Economics of Involuntary Resettlement: Questions and Challenges*. The World Bank, Washington DC.
- Cernea MM, McDowell C (Eds.) (2000) *Risks and Reconstruction: Experiences of Resettlers and Refugees*. The World Bank, Washington DC.
- Child BJ (2011) The absence of indigenous histories in Ken Burns's the national parks: America's best idea. *The Pub Hist* 33(2):24-29.
- Cochrane SMV, Matricardi EAT et al. (2017) Landsat-based analysis of mega dam flooding impacts in the Amazon compared to associated environmental impact assessments: Upper Madeira River example 2006–2015. *Remote Sensing Applications: Soc and Environ* 7:1-8. <https://doi.org/10.1016/j.rsase.2017.04.005>.
- Copeman V (1997) The impact of micro-hydropower on the aquatic environment. *Water Environ Jour* 11(6):431–435.
- Csiki S, Rhoads BL (2010) Hydraulic and geomorphological effects of run-of-river dams. *Prog in Phys Geog* 34(6):755-780. <https://doi.org/10.1177/0309133310369>.
- Csiki SJC, Rhoads BL (2014) Influence of four run-of-river dams on channel morphology and sediment characteristics in Illinois, USA. *Geomorphology* 206:215–229. doi:10.1016/j.geomorph.2013.10.0.
- Dak Psi (2023) *Dak Psi 3 and 4 Hydropower Project (2023)*. Project design document and other materials. Clean Development Mechanism, United Nations Framework Convention on Climate Change, Bonn, Germany. <https://cdm.unfccc.int/Projects/DB/KEMCO1307604296.67/view>.
- Dare GC, Murray RG et al. (2020) Run-of-river dams as a barrier to the movement of a stream-dwelling amphibian. *Ecosphere* 11(8):e03207.

- de Faria FAM, Jaramillo P (2015) Estimating greenhouse gas emissions from future Amazonian hydroelectric reservoirs. *Environ Res Lett* 10:art. 124019. <https://doi.org/10.1088/1748-9326/10/12/124019>.
- Dessaix J, Fruget JF et al. (1995) Changes of the macroinvertebrate communities in the dammed and by-passed sections of the French upper Rhone after regulation. *Reg Rivers: Res & Manag* 10(2-4):265-279.
- Doria CRC, Duponchelle F et al. (2018) Review of fisheries resource use and status in the Madeira River Basin (Brazil, Bolivia, and Peru) before hydroelectric dam completion. Taylor and Francis Inc., London.
- Du Toan (2022) Phát triển hài hòa thủy điện ở Kon Tum (Bài 2). <https://dantocmiennui.vn/phat-trien-hai-hoa-thuy-dien-o-kon-tum-bai-2-post324642.html>.
- Dyson M, Bergkamp G, Scanlon J (Eds.) (2003) *Flow: the essentials of environmental flows*. IUCN, Gland, Switzerland.
- Dépret T, Piégay H et al. (2019) Estimating and restoring bedload transport through a run-of-river reservoir. *Sci of the Total Environ* 654:1146–1157.
- EC 2022/2577. Council Regulation (EU) 2022/2577 of 22 December 2022 laying down a framework to accelerate the deployment of renewable energy. eur-lex.europa.eu/legal-content/DE/TXT/?uri=CELEX%3A32022R2577.
- Erlewein A (2014) The promotion of dams through the Clean Development Mechanism: Between sustainable climate protection and carbon colonialism. In *Large Dams in Asia: Contested Environments between Technological Hydroscares and Social Resistance*, 149-168.
- Fearnside PM (2006) Dams in the Amazon: Belo Monte and Brazil's Hydroelectric Development of the Xingu River Basin. *Environ Manag* 38(1):16-27. <https://doi.org/10.1007/s00267-005-00113-6>.
- Fearnside PM (2014a) Impacts of Brazil's Madeira River Dams: Unlearned lessons for hydroelectric development in Amazonia. *Environmental Sci & Policy* 38:164–172. <https://doi.org/10.1016/j.envsci.2013.11.004>.
- Fearnside PM (2014b) Brazil's Madeira River dams: a setback for environmental policy in Amazonian development. *Water Altern* 7(1),156-169. <https://bit.ly/3PoThwq>.
- Fearnside PM (2015) Tropical hydropower in the Clean Development Mechanism: Brazil's Santo Antônio Dam as an example of the need for change. *Clim Change* 131(4):575-589. <https://doi.org/10.1007/s10584-015-1393-3>.
- Fearnside PM (2017a) Planned disinformation: The example of the Belo Monte Dam as a source of greenhouse gases. In *Issberner L-R, Lena P (Eds), Brazil in the Anthropocene: Conflicts between Predatory Development and Environmental Policies*. Routledge, Taylor & Francis Group, New York, NY, 125-142. <https://doi.org/10.4324/9781315544069>.
- Fearnside PM (2017b) Brazil's Belo Monte Dam: Lessons of an Amazonian resource struggle. *Die Erde* 148(2-3):167-184. <https://doi.org/10.12854/erde-148-146>.
- Fearnside PM, Berenguer E et al. (2021) Drivers and impacts of changes in aquatic ecosystems. In: *Nobre C et al. (Eds), Amazon Assessment Report 2021*. Science Panel for the Amazon (SPA), UNSDSN, New York, USA. <https://doi.org/10.55161/IDMB5770>.
- Foran T, Manorom K (2009) Pak Mun Dam: Perpetually contested? In *Molle FF, Foran T, Käkönen M (Eds), Contested Waterscapes in the Mekong Region: Hydropower, Livelihoods and Governance*, 55-80. Earthscan, London.
- Freyhof J, Kottelat M (2008) Hucho. *The IUCN Red List of Threatened Species*. Version 8235.

- Fu X, Tao T et al (2008) Impacts of small hydropower plants on macroinvertebrate communities. *Acta Ecologica Sinica* 28(1):45–52.
- Gibeau P, Connors BM, Palen WJ (2017) Run-of-river hydropower and salmonids: potential effects and perspective on future research. *Can Jour of Fisheries and Aqua Sci* 74(7):1135-1149.
- Goldsmith E, Hilyard N (1984) *The Social and Environmental Effects of Large Dams*. Wadebridge Ecological Centre, Cornwall.
- Gonçalves AP (2011) *Ecologia e Etnoecologia de Hypancistrus zebra (Siluriformes: Loricariidae) no Rio Xingu, Amazônia Brasileira*. Master's dissertation in aquatic ecology and fisheries, Universidade Federal do Pará, Belém, PA, Brazil. 133 pp. <https://bityl.co/4XLC>.
- Goodland RJ, Juras A, Pachauri R (1993) Can hydro-reservoirs in tropical moist forests be environmentally sustainable? *Environ Cons* 20(2):122-130. <https://doi.org/10.1017/S0376892900037619>.
- Greimel F (2022) *Characterization of sub-daily flow fluctuations as a basis for sustainable hydropeaking management*. PhD thesis, University of Natural Resources and Life Sciences, Vienna, Austria.
- Greimel F, Zeiringer B et al. (2016) A method to detect and characterize sub-daily flow fluctuations. *Hydro Proc* 30:2063–2078. <https://doi.org/10.1002/hyp.10773>.
- Gunderson LH, Cosens B, Garmestani AS (2016) Adaptive governance of riverine and wetland ecosystem goods and services. *Journal of Environmental Management* 183: 353-360.
- Harlan T (2020) Conservation or decarbonization? Small hydropower and state logics of green development in China. *Ann of the Amer Assoc of Geog* 110(5):1464-1482. <https://doi.org/10.1080/24694452.2019.1684874>.
- Harrison JA (2019) Dams: test pros and cons case by case. *Nature* 566:315-317. <https://doi.org/10.1038/d41586-019-01137-2>.
- Hauser M, Doria CRC et al. (2019) Shedding light on the migratory patterns of the Amazonian goliath catfish, *Brachyplatystoma platynemum*, using otolith $^{87}\text{Sr}/^{86}\text{Sr}$ analyses. *Aqua Cons: Marine and Fresh Eco* 29:397–408.
- Haug J, Auer S et al. (2022) Retrofitting of existing bar racks with electrodes for fish protection—an experimental study assessing the effectiveness for a pilot site. *Water* 14(6):850.
- Hayes DS, Schülting L et al. (2022a) Hydropeaking: processes, effects, and mitigation. *Ref Mod in Earth Syst and Environ Sci* <https://doi.org/10.1016/B978-0-12-819166-8.00171-7>.
- Hayes DS, Schaufler G et al. (2022b) Hydro-morphological stressors reduce distribution range and affect population status of cyprinid fishes in Austria. *Front in Environ Sci* 10. <https://doi.org/10.3389/fenvs.2022.991722>.
- Hayes DS, Lautsch E et al. (2021) Response of European grayling, *Thymallus*, to multiple stressors in hydropeaking rivers. *Jour of Environ Manag* 292:112737. <https://doi.org/10.1016/j.jenvman.2021.112737>.
- Hayes DS, Brändle JM et al. (2018) Advancing towards functional environmental flows for temperate floodplain rivers. *Sci of the Total Environ* 633:1089-1104.
- Hennig T, Harlan T (2018) Shades of green energy: geographies of small hydropower in Yunnan, China and the challenges of over-development. *Glob Environ Chang* 49:116-128. <https://doi.org/10.1016/j.gloenvcha.2017.10.010>.
- Hirsch P (2016) The shifting regional geopolitics of Mekong dams. *Polit Geog* 51:63-74. <https://doi.org/10.1016/j.polgeo.2015.12.004>.

- Hocking MD, Faulkner SG (2021) Surprising salmonid response to water diversion at four run-of-river hydroelectric projects in British Columbia. *Can Jour of Fisheries and Aqua Sci* 78(10):1383-1396.
- Holčík J, Hensel K et al. (1988) The Eurasian Huchen, Hucho hucho, Largest Salmon of the World. Dr W. Junk Publisher, Dordrecht-Boston-Lancaster, pp. 239.
- Hunter MA (1992) Hydropower flow fluctuations and Salmonids: A review of the biological effects, mechanical causes and options for mitigation. State of Washington, Department of Fisheries, Olympia, WA.
- International Rivers Network (1999) An Update on the Environmental and Socio-Economic Impacts of the Nam Theun-Hinboun Hydroelectric Dam and Water Diversion Project in Central Laos. 15-17 August, 1999.
- Jager HI, Bevelhimer MS (2007) How run-of-river operation affects hydropower generation and value. *Environ Manag* 40(6):1004-15. <https://doi.org/10.1007/s00267-007-9008-z>.
- Jiang X, Lu D et al. (2018) Examining impacts of the Belo Monte hydroelectric dam construction on land-cover changes using multitemporal Landsat imagery. *App Geog* 97:35-47. <https://doi.org/10.1016/j.apgeog.2018.05.019>.
- Kahn JR, Freitas CE, Petrere M (2014) False shades of green: the case of Brazilian Amazonian hydropower. *Energies* 7(9):6063-6082. <https://doi.org/10.3390/en7096063>.
- Kelly-Richards S, Silber-Coats N et al. (2017). Governing the transition to renewable energy: A review of impacts and policy issues in the small hydropower boom. *Energy Policy* 101:251-264.
- Keppeler FW, Andrade MC (2022) Early impacts of the largest Amazonian hydropower project on fish communities. *Sci of the Total Enviro* 838:art. 155951. <https://doi.org/10.1016/j.scitotenv.2022.155951>.
- Kibler KM, Tullos DD (2013) Cumulative biophysical impact of small and large hydropower development in Nu River, China. *Water Res Res* 49:3104–3118. <https://doi.org/10.1002/wrcr.20243>.
- Kinew KA (1995) Manito Gitigaan. Governing in the Great Spirit's Garden. Wild Rice in Treaty #3. University of Manitoba, Winnipeg, MB.
- Knott J, Mueller M (2023a) Downstream fish passage at small-scale hydropower plants: Turbine or bypass? *Front in Environ Sci* 11:1168473.
- Knott J, Mueller M et al. (2023b) Ecological assessment of the world's first shaft hydropower plant. *Renew and Sust Energy Rev* 187:113727.
- Kraker D (2022) Did the historic flooding on Rainy Lake this summer have to be so bad? Minnesota Public Radio. <https://www.mprnews.org/story/2022/06/28/did-the-historic-flooding-on-rainy-lake-this-summer-have-to-be-so-bad>.
- Kuriqi A, Pinheiro AN et al. (2021) Ecological impacts of run-of-river hydropower plants—Current status and future prospects on the brink of energy transition. *Ren and Sust Energy Rev* 142:110833. <https://doi.org/10.1016/j.rser.2021.110833>.
- Li H, Zhao W et al. (2018) Entrainment effects of a small-scale diversion-type hydropower station on phytoplankton. *Ecol Eng* 116:45-51.
- Linares MS, Callisto M, Marques JC (2018) Thermodynamic based indicators illustrate how a run-of-river impoundment in neotropical savanna attracts invasive species and alters the benthic macroinvertebrate assemblages' complexity. *Ecol Indic* 88:181-189.

- Magalhães SB, da Cunha MC (Eds) (2017) A expulsão de Ribeirinhos em Belo Monte: Relatório da SBPC. Sociedade Brasileira para o Progresso da Ciência (SBPC), São Paulo, SP, Brazil, 448 pp. <http://portal.sbpnet.org.br/livro/belomonte.pdf>.
- Magilligan FJ, Roberts MO et al. (2021) The impact of run-of-river dams on sediment longitudinal connectivity and downstream channel equilibrium. *Geomorphology* 376:107568.
- Martins DEC, Seiffert MEB, Dziedzic M (2013) The importance of clean development mechanism for small hydro power plants. *Ren Energy* 60:643-647.
- McCully P (2001) *Silenced Rivers: The Ecology and Politics of Large Dams* (2nd edition). Zed Books, London.
- McManamay RA, Oigbokie CO et al. (2016) Classification of US hydropower dams by their modes of operation. *River Res and Applic* 32(7):1450-1468.
- McManamay RA, Peoples BK et al. (2015) Isolating causal pathways between flow and fish in the regulated river hierarchy. *Can Jour of Fisheries and Aqua Sci* 72:1731–1748. <https://doi.org/10.1139/cjfas-2015-0227>.
- Missingham BD (2003) *The Assembly of the Poor in Thailand: From Local Struggles to National Social Movement*. Silkworm Books, Chiang Mai.
- Moreira M, Hayes DS et al. (2019) Ecologically-based criteria for hydropeaking mitigation: a review. *Sci of the Total Environ* 657:1508–1522. <https://doi.org/10.1016/j.scitotenv.2018.12.107>.
- Neupane S, Das B et al. (2023) Impact of thirteen run-of-river hydroelectric projects on land use land cover and ecosystem services in Nepal. *Intern Jour of Energy and Water Res* 7(4):513-533.
- Pearson AJ, Pizzuto JE, Vargas R (2016) Influence of run of river dams on floodplain sediments and carbon dynamics. *Geoderma* 272:51-63.
- Pezzuti JCB, Zuanon J (2024) Brazil's Belo Monte license renewal and the need to recognize the immense impacts of dams in Amazonia. *Pers in Ecol and Cons* <https://doi.org/10.1016/j.pecon.2024.05.001>.
- Ponton DE, Lavoie RA et al. (2021) Understanding food web mercury accumulation through trophic transfer and carbon processing along a river affected by recent run-of-river dams. *Environ Sci & Tech* 55(5):2949-2959.
- Pracheil BM, DeRolph CR et al. (2016) A fish-eye view of riverine hydropower systems: the current understanding of the biological response to turbine passage. *Rev in Fish Biol and Fisheries* 26:153-167.
- Radinger J, van Treeck R, Wolter C (2022) Evident but context-dependent mortality of fish passing hydroelectric turbines. *Conserv Biol* 36(3):e13870.
- Randell H (2022) The challenges of dam-induced displacement: Reducing risks and rethinking hydropower. *One Earth* 5(8):849-852. <https://doi.org/10.1016/j.oneear.2022.07.002>.
- Reid P, Rajnovich G (1991) Laurel: a re-evaluation of the spatial, social and temporal paradigms. *Can Jour of Archaeo/Jour Can d'Archéolo*, 193-234.
- Ribas C, Muriel-Cunha J, Pezzuti JCB (2023) Devido à usina de Belo Monte, tragédia da seca já é realidade há tempos na Volta Grande do Xingu. *The Conversation*, 7 November 2023. <https://bit.ly/40VApMP>.
- Roberts TR (1995) Mekong mainstream hydropower dams: Run-of-the-River or ruin-of-the-river? *Nat Hist Bull Siam Soc* 43:9-19.
- Roberts TR (2001) On the river of no returns: Thailand's Pak Mun Dam and its fish ladder. *Nat Hist Bull Siam Soc* 49:189-230.

- Robson A (2013) Implications of small-scale run-of-river hydropower schemes on fish populations in Scottish streams. University of Hull, Hull.
- Rose V, Rollwagen-Bollens G et al. (2019) The effects of run-of-river dam spill on Columbia River microplankton. *River Res and Applic* 35(9):1478-1488.
- Sasges G, Ziegler AD (2023) We have eaten the rivers: The past, present, and unsustainable future of hydroelectricity in Vietnam. *Sustainability* 15(11):8969.
- Sasges G, Ziegler AD (2024) Problematic power: a perspective on the role of small hydropower in energy transitions in Vietnam. *ACS ES&T Water* 4(4):1242-1250. <https://doi.org/10.1021/acsestwater.3c00637>.
- Schmutz S, Jungwirth M et al. (2023) Der Huchen stirbt aus – was tun? Gefährdungsfaktoren und notwendige Maßnahmen in Bayern und Österreich. Sonderheft Österreichs Fischerei, Herausgegeben vom Österreichischen Fischereiverband, Wien.
- Schmutz S, Bakken TH et al. (2015) Response of fish communities to hydrological and morphological alterations in hydropeaking rivers of Austria. *River Res and Applic* 31:919–930. <https://doi.org/10.1002/rra.2795>.
- Schmutz S, Zitek A et al. (2002) Integrated approach to the conservation and restoration of Danube salmon, Hucho hucho, populations in Austria. In *Conservation of Freshwater Fishes: Options for the Future*, 157-173.
- Scudder T (2005) *The Future of Large Dams: Dealing with Social, Environmental, Institutional and Political Costs*. Earthscan, London.
- Sendzimi J, Schmutz S (2018) Challenges in riverine ecosystem management. In Schmutz S, Sendzimir J (Eds). *Riverine Ecosystem Management*, 1-16. https://doi.org/10.1007/978-3-319-73250-3_1
- Shoemaker B (1998) *Trouble on the Theun-Hinboun. A Field Report on the Socio-Economic and Environmental Effects of the Nam Theun-Hinboun Hydropower Project in Laos*. International Rivers Network, Berkeley, CA.
- Shoemaker B, Robichaud W (Eds) (2018) *Dead in the Water: Global Lessons from the World Bank's Model Hydropower Project in Laos*. University of Wisconsin Press, Madison, WI.
- Silva AT, Lucas MC et al. (2018) The future of fish passage science, engineering, and practice. *Fish and Fisheries* 19(2):340-362.
- Silverthorn VM, Bishop CA et al. (2018) Impact of flow diversion by run-of-river dams on American dipper diet and mercury exposure. *Environ Toxic and Chem* 37(2):411-426.
- Simonović P, Ristić R et al. (2021) Effects of run-of-river hydropower plants on fish communities in montane stream ecosystems in Serbia. *River Res and Applic* 37(5):722-731. <https://doi.org/10.1002/rra.3807>.
- Soukhaphon A, Baird IG (2024) Remembering the Mun: the role of women in engendering local geographies of resistance to the Pak Mun Dam. *Gender, Place and Culture* (published online). <https://doi.org/10.1080/0966369X.2024.1832713>.
- Sow MM, Majdi N et al. (2016) Retention of nutrients, suspended particulate matter and phytoplankton in a pondage associated with a run-of-the-river type hydroelectric power plant. *Ecology* 9(2):229-237. <https://doi.org/10.1002/eco.1633>.
- Strube J (2021) *Water, wild rice, and the ongoing production of Settler colonialism in Anishinaabe Akii*. The Pennsylvania State University.
- Strube J, Thomas KA (2021) Damming Rainy Lake and the ongoing production of hydro-colonialism in the US-Canada boundary waters. *Water Altern* 14(1):19-41.

- Sullivan Z (2016) Brazil's dispossessed: Belo Monte dam ruinous for Indigenous cultures. Mongabay, 8 December 2016. <https://bit.ly/41aHEAN>.
- Summer W, Stritzinger W, Zhang W (1994) The impact of run-of-river hydropower plants on temporal suspended sediment transport behaviour. *IAHS Publications-Series of Proc and Reports-Intern Assoc Hydro Sci* 224:411-420.
- Tashiro T, Kataoka T, Tsujimoto T (2015) Impacts of run-of-river weir water intakes on stream ecosystems: evaluation of ecosystem metabolism through continuous dissolved oxygen monitoring. *Jour of the Soc of Civil Eng, B1 (Hydrau Eng)* 71(4):I_1129-I_1134.
- Ullah EI, Ahmad S (2023) Hydrological and ecological impacts of run off river scheme; a case study of Ghazi Barotha hydropower project on Indus River, Pakistan. *Heliyon* 9(1):e09876. <https://doi.org/10.1016/j.heliyon.2023.e09876>.
- UNFCCC (2006) PROJECT DESIGN DOCUMENT FORM (CDM PDD) - Version 03, Dak Psi 3 and 4 Hydropower Project. United Nations Framework Convention on Climate Change (UNFCCC), New York. <https://cdm.unfccc.int/UserManagement/FileStorage/86HPNCWK1S0QFJOXI7LRGT5ZMBAVYE>.
- Venus TE, Hinzmann M et al. (2020) The public's perception of run-of-the-river hydropower across Europe. *Energy Policy* 140:111422. <https://doi.org/10.1016/j.enpol.2020.111422>.
- Von Sperling E (2012) Hydropower in Brazil: overview of positive and negative environmental aspects. *Energy Procedia* 18:110-118. <https://doi.org/10.1016/j.egypro.2012.05.023>.
- Waisberg LG (1984) An ethnographic and historical outline of the Rainy River Ojibway. In Noble, WC (Ed), *Historical Synthesis of the Manitou Mounds Site on the Rainy River*. Parks Canada, Cornwall, ON, 239-255.
- Wang H, Chen Y et al. (2016) Effects of the “run-of-river” hydro scheme on macroinvertebrate communities and habitat conditions in a Mountain River of Northeastern China. *Water* 8(1):31. <https://doi.org/10.3390/w8010031>.
- Wang Y, Wu N et al. (2022a) Small run-of-river hydropower dams and associated water regulation filter benthic diatom traits and affect functional diversity. *Sci of the Total Environ* 813:152566. <https://doi.org/10.1016/j.scitotenv.2021.152566>.
- Wang Y, Wu N et al. (2022b) Small run-of-river dams affect taxonomic and functional β -diversity, community assembly process of benthic diatoms. *Front in Ecol and Evol* 10:895328. <https://doi.org/10.3389/fevo.2022.895328>.
- Warren TJ (1999) A Monitoring Study to Assess the Localized Impacts Created by the Nam Theun-Hinboun Hydro-Scheme on Fisheries and Fish Populations. Theun-Hinboun Power Company (THPC), Vientiane.
- WCD (World Commission on Dams) (2000) *Dams and Development: A New Framework for Decision-Making*. Earthscan Publications, London.
- Whittington J. (2018) *Anthropogenic Rivers: The Production of Uncertainty in Lao Hydropower*. Cornell University Press, Ithaca, NY.
- Wildman LA, MacBroom JG (2005) The evolution of gravel bed channels after dam removal: Case study of the Anaconda and Union City Dam removals. *Geomorphology* 71(1-2):245-262. <https://doi.org/10.1016/j.geomorph.2005.02.008>.
- Wu N, Tang T et al. (2010) Impacts of cascade run-of-river dams on benthic diatoms in the Xiangxi River, China. *Aqua. Sci.* 72:117–125. <https://doi.org/10.1007/s00027-009-0121-3>.
- Wu N, Tang T et al. (2009) Changes in benthic algal communities following construction of a run-of-river dam. *Jour Am Benthol Soc* 28(1):69–79. <https://doi.org/10.1899/08-047.1>.

- Zanon FM, Pineda A et al. (2024) Impact of a run-of-river dam in an Amazonian large river on the spatial and temporal patterns of phytoplankton beta diversity. *Bio Cons* 295:110656. <https://doi.org/10.1016/j.biocon.2024.110656>.
- Zengin M, Kurtoğlu O et al. (2017) Impact of run-of-river hydropower plants operation on aquatic ecosystem and trout (*Salmo labrax*) population in the eastern Black Sea region. *Aqua Sci and Eng* 32(4):189-207. <https://doi.org/10.31238/ase.377>.
- Zhou S, Tang T et al. (2008) Impacts of a small dam on riverine zooplankton. *Intern. Rev Hydrobiol* 93(3):297–311. <https://doi.org/10.1002/iroh.200711038>.
- Ziv G, Baran E et al. (2012) Trading-off fish biodiversity, food security, and hydropower in the Mekong River Basin. *Proc of the Nat Acad of Sci* 109:5609–5614. <https://doi.org/10.1073/pnas.1201423109>.

Acknowledgements & Funding

Alan Ziegler was supported from funding support from the National Science, Research and Innovation Fund (NSRF) via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation (grant number B48G660115). This research was also funded in part by the Austrian Science Fund (FWF) [P 34061-B].